

COVID-19 and the Mental Health of Adolescents in British Columbia

November 2023

Jane Friesen, Brian Krauth and Brett Wilmer¹

Abstract

We use linked individual-level data on school enrollment, physician services received, and prescription medications from September 2016 to March 2022 to measure the effect of the COVID-19 pandemic and associated disruptions on the monthly frequency of mental health treatments received by adolescents in British Columbia. The results suggest substantial increases for non-Indigenous English home language girls in treatment for depression/anxiety, ADHD, eating disorders and other mental health conditions. Non-English home language girls also show increases in treatment for depression/anxiety and Indigenous girls show increases in treatment for ADHD. In contrast, boys show no change or even reductions in treatment for most mental health conditions. The exception is conduct disorders, which increase among English home language boys. Treatment for substance use disorders declines among English home language boys, Indigenous boys and Indigenous girls. Effects vary somewhat by socioeconomic status, but we find no evidence that they vary substantially by schooling mode.

¹ Jane Friesen, Department of Economics, Simon Fraser University, Burnaby BC Canada, friesen@sfu.ca; Brian Krauth, Department of Economics, Simon Fraser University, Burnaby BC Canada, bkrauth@sfu.ca; Brett Wilmer, Director, Provincial Statistics, British Columbia Ministry of Citizen Services, Victoria BC Canada, brett.wilmer@gov.bc.ca. Lindsay Forestell provided excellent research assistance. All inferences, opinions, and conclusions drawn in this paper are those of the authors, and do not reflect the opinions or policies of the Data Innovation Program or the Province of British Columbia. The following material was developed as part of *Assessing the Effect of COVID-19 on Mental Health Disorders among B.C.'s K-12 Student Population* project, commissioned by the Ministry of Education and Childcare, Province of British Columbia (SFU Research Ethics # 30000383). Latest version available at <https://bvkrauth.github.io/publication/covidmh>.

1 Introduction

Isolation from face-to-face contacts, reduced access to school-based mental health and support services, limits on opportunities for physical activity and changes in exposure to home environments brought on by the global COVID-19 pandemic may have seriously undermined the mental health of children and youth. Poor mental health among children and youth is of direct concern because of the suffering that it entails and because even short-term mental health problems can have serious consequences in childhood and adolescence (Busch et al. 2014; Currie and Stabile 2009). Moreover, a majority of mental health disorders in adulthood begin before the age of 15 (Kim-Cohen et al. 2003; Kessler et al. 2001; 2007) and the effects of conditions like anxiety and depression on decision-making, risk-taking and preferences (Ridley et al. 2020) may alter students' pathways through school and work and are associated with undesirable adult outcomes including reduced earnings and well-being (Prinz et al. 2018).

This paper investigates the use of physician services and prescription medications to treat mental health disorders among adolescents before and during the COVID-19 pandemic. We use longitudinal student-level education records for the Grade 8-11 school-age population in British Columbia (B.C.), linked to physicians' billing records from the province's universal public health insurance system and the universe of community pharmacy prescription medication dispensing events from September 2016 to March 2022. We estimate before-after and event study models for monthly physician services related to depression and anxiety, attention-deficit/hyperactivity disorders (ADHD), psychotic disorders, eating disorders, substance use disorders and conduct disorders, and for three categories of psychotropic medications commonly prescribed to treat mental health conditions in adolescents – antidepressant/anti-anxiety medications, stimulants and other medications commonly used for management of ADHD symptoms, and anti-psychotics.

Our work extends the small but growing literature that uses population health records to track mental health-related interactions with the health care system among adolescents during the pandemic. Previous studies that use population health records to study changes in mental health treatment during the pandemic have investigated physician services related to overall mental health conditions, mood disorders, psychotic disorders, substance use disorders and neurodevelopmental problems in Ontario (Saunders et al. 2022; Toulany et al. 2023), overall mental health care treatment (Hvide and Johnsen 2022), ADHD, anxiety, depression and sleep disorders (Evensen et al. 2022), and eating disorders (Surén et al. 2022) in Norway, eating disorders, autism, ADHD, panic disorders, hypochondriasis, anxiety disorders, anorexia nervosa, depressive disorders and bipolar disorders in Korea (Kim et al. 2022), and prescriptions for antidepressant medication in Australia (Costa et al. 2022) and antidepressant, anxiolytic/sedative-hypnotic, antipsychotic and stimulant medications in Manitoba (Leong et al. 2022). In

general, these studies have found higher than predicted levels of mental health care interactions beginning either at the onset of the pandemic or following periods of lock-down and continuing through the end of 2020 and into 2021. Studies that provide separate analysis for males and females find little (Kim et al. 2022; Evensen et al. 2022; Saunders et al. 2022) or no (Costa et al. 2022) effect of the pandemic on physician services related to mental health among adolescent boys, but significant increases in some cases among girls. Toulany et al. (2023) find that mental health-related physician visits by children and youth in Ontario during the first year of the pandemic were higher than predicted only among immigrants (but not among non-immigrants or refugees), lower than predicted among those living in low SES neighborhoods and higher than predicted among those living in high SES neighborhoods.

We extend this literature in several ways. First, our data extend to March 2022, allowing us to investigate the extent to which any changes in the use of mental health services during the first year of the pandemic persisted through the following year. Second, in light of evidence that mental health care use both varied across groups defined by ethnicity and immigrant status² and responded differently during the pandemic (Toulany et al. 2023), we analyze mental health trends separately for Indigenous students and for two groups of non-Indigenous students that we identify in the education records: those who speak English at home and those who speak a different language at home. While home language is likely to be correlated with immigrant status, our definition of groups by Indigeneity and home language is very different from the strict measures of children's immigrant status used by Toulany et al. (2023). We are not aware of any previous studies that provide specific analysis of Indigenous youth in this context. Third, differences in learning modes adopted by school districts when lock downs ended may have implications for the socialization and stress that adolescents encountered during the pandemic. Compared to full-time in-person instruction, hybrid learning (i.e., a combination of in-person and remote (online) instruction) may involve loss of regular routine, reduced social contact with peers (for better or for worse) and increased demands on self-motivation and self-regulation.^{3 4} Following a period of school closure in spring of 2020,

² Gadermann et al. (2022) find that first-generation immigrant and refugee children in British Columbia had the lowest mental health diagnostic prevalence rates during the pre-COVID period, followed by second-generation immigrants and refugees. A large body of evidence shows that native-born Caucasians in high income countries are much more likely to access the mental health care system than other groups, with the lowest prevalence rates among self-identified Asians and first-generation immigrants (e.g., Abe-Kim et al. 2007; Cook et al. 2017; Georgiades et al. 2018). Clinically significant cases of mental illness may go undiagnosed when parents, teachers or physicians fail to recognize children's needs or because of constraints on information, time, or availability of care, or because of stigma or discrimination (Bharadwaj et al. 2017; Thornicroft et al. 2017).

³ Previous research finds that both adolescents diagnosed with ADHD and their parents have reported students having difficulty engaging in remote learning during the pandemic (Becker et al. 2020; Sibley et al. 2021).

⁴ Using cross-sectional data from a December 2020 survey in the U.S., Hawrilenko et al. (2021) find that parent-reported difficulties were higher among children and adolescents enrolled in hybrid programs compared to in-person programs and were similar to students in remote schooling. Svaleryd et al. (2021) compare the change in services of psychiatric specialists and prescriptions for psychotropic medication during COVID among students at Swedish upper secondary schools, which closed for three months at the onset of the pandemic, reopened full-time in the fall,

most B.C. students returned to full-time in-person learning in September, but many school districts implemented hybrid learning models in some high school grades. We use this variation to implement an estimator that isolates the effect of hybrid versus face-to-face programs under the maintained assumption that underlying trends and seasonal patterns are the same across grades and between districts that offered hybrid versus face-to-face instruction.

Like previous authors, we find strikingly different results for adolescent girls versus boys. Following a period of substantial decline in the early months of the pandemic among both sexes, physician services and prescription medications related to mental health disorders overall recovered among boys but did not deviate substantially from predicted levels thereafter. The picture for adolescent girls is very different – physician services and prescription medications recovered to normal levels within the first half-year of the pandemic and then rose substantially and remained high relative to previous levels through to the end of our data period in March 2022.

Our detailed investigation reveals distinct results by demographic group and diagnostic category. Among boys, we see a statistically significant increase in physician visits only in the case of conduct disorders among those from English home language families, and no cases where prescriptions increased. Notably, physician visits related to substance use disorders *fell* substantially among English home language boys (-31.1%), Indigenous boys (-43.3%) and Indigenous girls (-27.7%).

Physician visits related to ADHD increased substantially among Indigenous girls (32.3%). Among non-Indigenous girls, we find striking and precisely estimated increases among those from English home language families in physician visits related to depression and anxiety (13.1%), ADHD (28.3%), psychotic disorders (17.5%), conduct disorders (55.6%) and eating disorders (48.4%), in prescription medications used to treat depression and anxiety (5.1%) and ADHD (7.6%), and for anti-psychotics (18.4%). Among non-English home language girls, physician visits related to depression and anxiety increased substantially (22.1%). In all of these cases, the increase in treatment is sustained at or near its post-COVID peak into early 2022, when our data end.

Further investigation shows that increases in physician services were greater among high SES girls in cases where high SES girls were already more likely to receive treatment in the pre-COVID period, e.g., eating disorders among English home language girls and depression and anxiety among non-English

and moved to hybrid classes from January through March of 2021, to those of students at lower secondary schools, which remained open full-time throughout the period. They find that specialist mental health contacts and medication use fell among both groups relative to trend, but more so among students who experienced school closures. This relative decline persisted into 2021, when upper secondary schools were offering hybrid learning.

home language girls. We also find that the tendency for low-SES girls to be disproportionately prescribed anti-psychotics increased during the pandemic.

Finally, we find no evidence that the increases in physician services and prescriptions that we observe are different for students who attended hybrid versus face-to-face school programs.

2 Data sources and restrictions

We use student-level administrative data from annual sets of three linked files: school enrolment records maintained by the B.C. Ministry of Education and Childcare,⁵ records of physician billing under B.C.'s universal Medical Services Plan (MSP),⁶ and prescription medication dispensing records maintained under B.C.'s Pharmanet system.⁷ We accessed anonymized extracts of these data files via Population Data B.C.'s Secure Research Environment (<https://www.popdata.bc.ca>). The extracts we use here include Ministry of Education records from the 2016/17 through 2021/22 school years and MSP and Pharmanet records from September 2016 through March 2022. We link student-level records across files using a unique Study ID that is included in each file. To minimize residual disclosure risks, all sample sizes are reported rounded to the nearest 100.

The Ministry of Education⁸ file includes a small set of demographic characteristics – birth month and year, gender, language spoken at home, British Columbia residency, Aboriginal self-identification (for which we use the term Indigenous), First Nations on-reserve residency, district and school attended, grade, and program information (i.e., Special Needs, English as a Second Language and French Immersion) for each year the student is enrolled in a private or public school in B.C. The Ministry also provides a student-level index of neighborhood socioeconomic status, constructed from Census characteristics at the Dissemination Area (DA) level and linked to individual students by postal code of residence.⁹

All B.C. residents are required to enroll in MSP, which covers the cost of all medically necessary services provided or ordered by a registered physician. Each insured individual has a unique Personal Health

⁵ British Columbia Ministry of Education. K-12 Student Demographics and Achievements V03. Data Innovation Program, Province of British Columbia. Data Extract. 2022.

⁶ British Columbia Ministry of Health. Medical Services Plan (MSP) Payment Information File V03. Data Innovation Program, Province of British Columbia. Data Extract. 2022.

⁷ British Columbia Ministry of Health. Pharmanet V02. Data Innovation Program, Province of British Columbia. Data Extract. 2022.

⁸ The Ministry of Education and Childcare was known as the Ministry of Education during the time period covered by our data.

⁹ British Columbia Ministry of Education. K to 12 Socio Economic (SES) Index V01. Data Innovation Program, Province of British Columbia. Data Extract. 2022. For details of the construction of this variable see “Metadata for K to 12 Socio Economic Status (SES) Index - E03 - Datasets - Data Catalogue (gov.bc.ca)” at <https://catalogue.data.gov.bc.ca/dataset/htdd>.

Number that is used for billing purposes (Government of British Columbia 2022a). The MSP file used in this study includes a record for each insured service provided to a covered individual, except for those related to workplace injuries that are covered by the B.C. Workers' Compensation Program. Physicians may bill outside the MSP system but may not charge more than the MSP prescribed rate for services; patients who pay these doctors directly may apply for reimbursement at the MSP rate (Government of British Columbia 2022b). Physicians who are reimbursed under an alternative payment plan (i.e., not fee-for-service) are required to submit the same documentation as fee-for-service physicians. All of these services are captured in our data. The MSP records include the date of service, specialty of the physician delivering the service, a billing code for the type of service provided, a diagnostic code based on the ICD-9 classification system, delivery location and additional information related to billing.

Pharmanet is a province-wide network that, since 1995, has linked all B.C. community (as opposed to hospital or institution-based) pharmacies to a central data system. Every prescription medication dispensed by a community pharmacy in B.C. must be recorded on the Pharmanet system. Prescription medications dispensed in hospitals or residential treatment facilities are not captured in the Pharmanet records (Government of British Columbia 2022b). The Pharmanet records include the date the prescription was dispensed, the drug name and drug identification number (DIN), the dosage and the number of days supplied.

Our primary outcome variables are indicators of whether a student received mental health-related physician services (in the MSP data) or filled a mental health-related prescription (in the Pharmanet data) during a given month or academic (September to August) year. We define a physician service reported in the MSP data as mental health-related if the reported ICD-9 diagnostic code is associated with "organic psychotic conditions", "other psychoses", or "neurotic disorders, personality disorders and other non-psychotic mental disorders" or if the reported Ministry of Health code is associated with "Anxiety/depression". See Appendix Table A1 for details. We identify mental health-related prescription medications using a list of DIN codes associated with all antidepressant, anti-psychotic, sedative and anxiolytic drugs that were dispensed to children aged 5-19 in B.C. between 2009/10 and 2020/21. This list was prepared by the B.C. Ministry of Health from Pharmanet data and drug categories. To this list we add medications approved for use in Canada for the treatment of ADHD in children. See Appendix Table A2 for details.

An important limitation of our data is that we do not observe mental health services provided by psychologists, counsellors, or social workers, whose services are not covered by MSP. However, any course of treatment requiring prescription medication will involve a physician, and both family physicians

and psychiatrists often play a role in both the initial diagnosis and subsequent management of mental health conditions.

3 Pre-COVID patterns

We begin our analysis by identifying cross-sectional relationships among key variables during a period that predates the beginning of the COVID pandemic in early 2020. These patterns and relationships are of direct interest and will also inform our methodology for measuring the impact of COVID.

Table 1 presents descriptive statistics for key student characteristics (measured annually) during the pre-COVID period, from the beginning of the 2016/17 (September to August) school year to the end of the 2018/19 school year, as well as the cross-sectional relationship between those characteristics and our annual measures of mental health care interactions. The first column provides population shares for each characteristic. Our grade classification is based on age-appropriate grade – the grade normally observed for the student’s birth year – since some secondary students do not have an assigned grade. Gender is reported as male or female in the enrollment data. Our measures of language and cultural identity are based on a constructed variable in the enrollment data for whether a student has ever self-identified as Indigenous, and on a variable reporting the language spoken in the student’s home. As one would expect, students in our data are roughly evenly distributed by age-appropriate grade and gender. Just over 12% of students self-identify as Indigenous and just over 63% of students live in non-Indigenous families that speak English at home. The next largest home language group is Chinese (6.5% of all students), followed by Punjabi (5.4%). The remaining 12.7% of students report one of dozens of other home languages.

Table 1: Student characteristics and mental health treatment received, pre-COVID (2016/17 through 2018/19).

Sub-population	% of total	Annual % receiving treatment	
		Physician service	Prescription filled
All students	100.0	14.4	9.2
Grade (age)			
8 (age 13)	25.0	11.8	6.9
9 (age 14)	25.0	13.8	8.4
10 (age 15)	25.0	15.4	10.0
11 (age 16)	25.0	16.8	11.6
Gender			
Female	48.6	16.3	10.2
Male	51.4	12.7	8.3
Language/cultural identity			
Indigenous	12.3	22.5	15.4
Non-Indigenous by home language			
English	63.2	15.7	10.4
Chinese	6.5	6.1	2.3
Punjabi	5.4	6.8	2.6
Other	12.7	7.7	3.5

Population consists of annual observations on B.C. resident students ages 13-16 on December 31 of the current academic year ($n \approx 552,400$ student-years). Annual percent receiving treatment is based on mental health-related services received or prescriptions filled between September and August of that academic year. See Appendix A for definitions.

The remaining columns in Table 1 report the average share of students for whom at least one mental health-related physician service was billed to MSP during a given school year and the average share who filled at least one mental health-related prescription. On average, 14.4% of students received a mental health-related physician service at least once in a given year. A smaller share (9.2%) filled a mental health-related prescription. The share receiving a mental health-related physician service is higher among older students, rising from 11.8% for those in the grade 8 age group to 16.8% in grade 11, while the share filling a mental health-related prescription rises from 6.9% in the grade 8 age group to 11.6% in grade 11. Gender and cultural background both play large roles in whether students receive mental health treatment. The share of students receiving a mental health-related physician service was greater among girls than boys (16.3% of girls versus 12.7% of boys), as was the share filling a mental health-related prescription (10.2% of girls versus 8.3% of boys). Indigenous students in our data are substantially more likely than non-Indigenous English home language students to receive a mental health-related physician service (22.5% versus 15.7%) and to fill a mental health-related prescription (15.4% versus 10.4%). The share of students receiving mental health treatment is strikingly lower among those from non-Indigenous families that do not speak English at home. For example, only 6.1% of students from Chinese home language families and 6.8% of students from Punjabi home language families receive a mental health-related physician service, compared to 15.7% of students from English home language families. These differences are even greater for mental health-related prescriptions: for example, only 2.3% of Chinese home language students and 2.6% of Punjabi home language students, compared to 10.4% of English home language students.

We further characterize the pre-COVID baseline by disaggregating the physician service data according to the diagnostic category associated with each record. Table 2a shows that the most frequently observed mental health diagnostic category is depression and anxiety (overall, 10% of students have at least one physician record with this diagnostic code in an average year), followed by ADHD (3.7%), psychotic disorders (1.9%), conduct disorders (0.7%), substance use disorders (0.7%) and eating disorders (0.3%). Table 2b shows that prescription rates were highest for antidepressants/anti-anxiety medications (6.4%), followed by medications commonly used to treat ADHD (3.6%) and anti-psychotics (1.3%). Note that there is not a simple mapping between medications and diagnostic categories, as a given medication may be used to treat multiple conditions and multiple medications may be used to treat a given condition. Treatment for all categories of mental health disorders is more frequent in higher grades in most cases, with ADHD and conduct disorders being notable exceptions where treatment is less frequent among older students.

Gender and cultural background also have differential impacts across diagnostic categories and groups of medications. Girls are almost twice as likely as boys to receive a physician service related to a diagnosis of depression/anxiety (13.2% of girls versus 6.9% of boys) or to fill a prescription for an antidepressant or anti-anxiety medication (8.7% of girls versus 4.3% of boys) and are more than six times as likely to receive a physician service related to an eating disorder (0.6% of girls versus <0.1% of boys). Boys are more than twice as likely as girls to receive a physician service related to ADHD (5.1% of boys versus 2.2% of girls) or to fill a prescription for an ADHD-related medication (4.8% of boys versus 2.2% of girls). Indigenous students have the highest frequencies of physician services related to all categories of mental health conditions except eating disorders. Among non-Indigenous students, English home language students are at least twice as likely to receive a mental health-related physician service in every diagnostic category and are more than three times as likely to fill a mental health-related prescription as those who speak another language at home.

Table 2a: Share of students receiving mental health-related physician services in a given school year by condition, pre-COVID (2016/17 through 2018/19).

Sub-population	Annual % receiving physician services						
	Depression/ Anxiety	ADHD	Psychosis	Conduct	Substance	Eating	Other
All students	10.0	3.7	1.9	0.7	0.7	0.3	1.8
Grade (age)							
8 (age 13)	7.0	4.1	1.5	0.8	0.3	0.2	1.4
9 (age 14)	9.2	3.9	1.8	0.7	0.6	0.3	1.7
10 (age 15)	11.0	3.5	2.1	0.6	0.8	0.4	1.9
11 (age 16)	12.6	3.2	2.2	0.6	1.0	0.4	2.1
Gender							
Female	13.2	2.2	2.0	0.5	0.7	0.6	2.2
Male	6.9	5.1	1.9	0.8	0.6	<0.1	1.4
Language/cultural identity							
Indigenous	15.5	6.0	3.3	1.4	2.1	0.3	2.8
Non-Indigenous by home language							
English	10.9	4.1	2.1	0.7	0.6	0.4	1.8
Chinese	4.2	1.0	1.0	0.2	0.1	0.2	1.0
Punjabi	4.6	1.0	0.6	0.4	0.3	0.1	1.2
Other	5.2	1.6	1.0	0.3	0.3	0.2	1.2

Population consists of annual observations of B.C. resident students ages 13-16 on December 31 of each academic year ($n \approx 552,400$ student-years). Annual percent receiving physician services is based on mental health-related physician services received between September and August of that academic year. See Appendix A for definitions of diagnostic categories.

Table 2b: Share of students filling mental health-related prescriptions in a given school year by drug category, pre-COVID (2016/17 through 2018/19).

Sub-population	Annual % filling prescriptions		
	Antidepressant and related	ADHD-related	Antipsychotic
All students	6.4	3.6	1.3
Grade (age)			
8 (age 13)	3.7	3.9	0.9
9 (age 14)	5.5	3.7	1.1
10 (age 15)	7.3	3.5	1.4
11 (age 16)	9.2	3.3	1.7
Gender			
Female	8.7	2.2	1.4
Male	4.3	4.8	1.2
Language/cultural identity			
Indigenous	10.7	6.1	2.7
Non-Indigenous by home language			
English	7.3	4.1	1.3
Chinese	1.5	0.8	0.3
Punjabi	1.9	0.7	0.3
Other	2.5	1.2	0.5

Population consists of annual observations of B.C. resident students ages 13-16 on December 31 of current academic year ($n \approx 552,400$ student-years). Annual percent filling prescriptions is based on mental health-related prescriptions filled between September and August of that academic year. See Appendix A for definitions of prescription categories.


The clear heterogeneity by gender and cultural background in rates of mental health treatment suggests that COVID impacts on treatment may exhibit similar heterogeneity. We therefore conduct our main analysis by subpopulations defined by gender and cultural background (pooling Chinese, Punjabi and other home language students). In addition, depression/anxiety and ADHD account for a large proportion of mental health treatment among adolescents, and gender and age patterns differ substantially for these two conditions. For this reason, our main analysis treats each broad diagnostic and drug category separately.

4 COVID impacts

In order to investigate patterns of mental health care interactions during the pandemic, we turn to monthly indicators of physician visits and prescriptions filled. Changes in these monthly measures will reflect both changes on the extensive margin (i.e., the number of students entering into and exiting treatment) and changes on the intensive margin (i.e., the frequency of services among those who are in treatment). The time series plot of the monthly share of students receiving mental health-related physician services through March 2022 in Figure 1a shows several clear patterns. During the pre-COVID period, the series shows an upward trend over time and a repeating monthly pattern, with the share receiving physician services typically rising over the school year to a May peak, followed by a substantial decline during the summer months. This pattern then restarts in September as each cohort of students progresses to the next grade. The predicted values from a regression of this series on monthly indicators and a linear trend

(estimated from data before March 2020) shown alongside the raw data in Figure 1a have a close within-sample fit to the actual values but deviate substantially from their predicted values during the COVID period. Instead of the usual seasonal rise in spring of 2020, mental health-related physician services declined during these early COVID months. Physician services returned to normal seasonal rates at the end of the summer followed by a steep and sustained increase relative to predicted values through March 2022, when our data series ends. These dynamics are likely driven by a combination of social factors influencing mental health and changes in access to mental health care. The initial decline is at least partly attributable to service interruptions for all forms of health care in the early months of COVID, and the subsequent increase may be partly attributable to “catch-up” as individuals who would have sought treatment earlier are finally able to access care. Our more detailed analysis below will consider both the net effect and the month-by-month dynamics.

Figure 1a. Actual and predicted monthly share receiving mental health-related physician services, September 2016 to March 2022. Predicted values are based on regression fit to observations before March 2020.




Corresponding plots for the share of students filling a mental health-related prescription in Figure 1b show similar patterns.¹⁰ Mental health-related prescriptions were slightly above predicted rates in January and February 2020, followed by a pronounced spike in March 2020 that suggests that individuals may have been filling prescriptions in anticipation of pandemic-related disruptions. Prescriptions fell below

¹⁰ The mental health physician services and mental health prescriptions series track one another quite closely, with more students filling a prescription than receiving physician services in a given month. Similar patterns can be observed when we disaggregate by diagnostic or drug category, though the steepness of the trend and timing of the seasonal peaks varies somewhat across categories.

predicted rates through the summer and then began to rise, exceeding predicted rates by late fall and remaining above-trend through March 2022.

Figure 1b. Actual and predicted monthly share filling mental health-related prescriptions, September 2016 to March 2022. Predicted values are based on regression fit to observations before March 2020.



Figures 2a and 2b present similar monthly time series by gender. It is immediately apparent that both the increase in physician services and in prescriptions seen in the aggregate data are driven almost exclusively by substantial increases among girls. Among boys, physician services rose back to their predicted level by September 2020, with some evidence of a relatively small increase relative to predicted levels emerging among boys in the second half of the 2020/21 school year and re-emerging during the 2021/22 school year. Mental health-related prescriptions did not return to their predicted level among boys until the summer of 2021.

Figure 2a. Actual and predicted monthly share receiving mental health-related physician services, by gender, September 2016 to March 2022. Predicted values are based on regression fit to observations before March 2020.





Figure 2b. Actual and predicted monthly share filling mental health-related prescriptions, by gender, September 2016 to March 2022. Predicted values are based on regression fit to observations before March 2020.



4.1 Main results

We next use a simple regression model to quantify the difference between the monthly average of each outcome variable between March 2020 and March 2022 and its pre-COVID average, net of seasonal effects and a linear trend and conditional on covariates. This measure reveals whether the magnitude and

duration of any positive deviations from predicted outcomes in later months were sufficient to compensate for the negative deviations observed during the early months of the pandemic. It provides a simple summary measure of the “COVID effect”, which we interpret as inclusive of the effect of changes both to the social environment and to health care access. We estimate this effect using a simple before/after regression model:

$$y_{it} = \beta \times 1(e_t \geq 0) + \mu_{m(t)} + \lambda t + \alpha X_{it} + \epsilon_{it} \quad (1)$$

where y_{it} is an indicator that student i received a mental health treatment during month t , e_t indicates event time relative to March 2020, $\mu_{m(t)}$ is a set of month fixed effects, λt is a monthly time trend, and X_{it} is a set of student-level covariates that includes age-appropriate grade (i.e. the grade the student would normally be enrolled in based on date of birth), neighborhood (i.e., Census Dissemination Area) SES quartile and birth quarter.¹¹ The parameter β captures the effect of interest.

Table 3a: COVID effect on mental health-related physician services by sub-population, as percentage change relative to pre-COVID baseline.

Sub-population	COVID effect on monthly share receiving physician services (as % of pre-COVID mean)					
	Depression/ Anxiety	ADHD	Psychosis	Conduct	Substance	Eating
Girls						
Indigenous ($n \approx 773,400$)	3.1 (3.1)	32.3*** (7.1)	0.3 (7.9)	19.5 (14.5)	-27.7** (11.0)	37.8 (25.2)
Non-indigenous by home language						
English ($n \approx 3,863,000$)	13.1*** (1.8)	28.3*** (4.8)	17.5*** (5.0)	55.6*** (11.1)	1.4 (9.9)	48.4*** (9.2)
Non-English ($n \approx 1,466,800$)	22.1*** (4.8)	12.8 (13.6)	3.7 (13.0)	8.7 (30.4)	-30.0 (30.0)	20.0 (18.9)
Boys						
Indigenous ($n \approx 788,300$)	-11.1** (4.7)	1.4 (4.9)	-2.1 (9.5)	18.5 (11.7)	-43.3*** (13.9)	
Non-indigenous by home language						
English ($n \approx 4,024,200$)	-6.3*** (2.4)	0.4 (2.5)	-0.8 (4.6)	19.1*** (6.6)	-31.1*** (11.5)	
Non-English ($n \approx 1,631,800$)	6.4 (6.1)	2.5 (7.8)	5.7 (11.4)	10.9 (17.4)	20.0 (24.0)	

Table reports coefficient on post-COVID indicator expressed as a percentage change in the pre-COVID baseline; control variables include time trend and indicators for month, grade, neighborhood SES quartile, and birth quarter. Estimation sample includes observations from September 2016 through March 2022, with reported sample size based on number of unique student-month pairs and rounded to nearest 100. Eating disorder results are not reported for boys due to limited variation in the dependent variable. Standard errors clustered by student in parentheses. *** = 1%, ** = 5%, * = 10%.

¹¹ The inclusion of birth quarter as a control is motivated by the well-documented relative age effect in ADHD diagnoses (Evans et al. 2010; Furzer et al. 2022), and by evidence that being younger in grade is associated with reduced life satisfaction, lower self-esteem, reduced confidence in abilities, peer problems and increased internalizing symptoms (Thompson et al. 2004; Crawford et al. 2011; Patalay et al. 2015; Fumarco and Baert 2019). We find that none of the relative age effects in our regression models are statistically significant.

Tables 3a and 3b show our main before/after results, where we report effect sizes as percentage changes and standard errors are clustered by student.¹² The corresponding regression coefficient estimates from equation (1) are reported in Appendix Tables B2a and B2b, and baseline frequencies used to convert these estimates into percentage terms are reported in Appendix Tables B1a and B1b.

Table 3b: COVID effect on mental health-related prescriptions filled by sub-population, as percentage change relative to pre-COVID baseline.

Sub-population	COVID effect on monthly share filling prescriptions (as % of pre-COVID mean)		
	Antidepressant and related	ADHD-related	Antipsychotic
Girls			
Indigenous (<i>n</i> ≈ 773,400)	0.3 (3.5)	7.1 (5.3)	3.5 (8.3)
Non-indigenous by home language			
English (<i>n</i> ≈ 3,863,000)	5.1*** (1.9)	7.6** (3.6)	18.4*** (5.0)
Non-English (<i>n</i> ≈ 1,466,800)	2.2 (6.5)	-26.0** (12.7)	24.1 (15.8)
Boys			
Indigenous (<i>n</i> ≈ 788,300)	-6.7 (5.1)	-4.9 (3.8)	5.4 (6.7)
Non-indigenous by home language			
English (<i>n</i> ≈ 4,024,200)	-5.0** (2.5)	-8.6*** (2.1)	1.6 (4.7)
Non-English (<i>n</i> ≈ 1,631,800)	6.7 (8.3)	-15.8** (7.5)	-17.2 (14.2)

Table reports coefficient on post-COVID indicator expressed as a percentage change in the pre-COVID baseline; control variables include time trend and indicators for month, grade, neighborhood SES quartile and birth quarter. Estimation sample includes observations from September 2016 through March 2022, with reported sample size based on number of unique student-month pairs and rounded to nearest 100. Standard errors clustered by student in parentheses. *** = 1%, ** = 5%, * = 10%.

We highlight five key results. First, physician services related to ADHD increased substantially among Indigenous girls (32.3%). Second, among English home language girls, physician services increased for depression and anxiety disorders (13.1%), ADHD (28.3%), psychotic disorders (17.5%), conduct disorders (55.6%) and eating disorders (48.4%), with smaller increases in prescription medications used to treat depression and anxiety (5.1%) and ADHD (7.6%) and for anti-psychotics (18.4%). Note that depression/anxiety is by far the most common mental health diagnosis for this age group, so even a moderately-sized percentage increase affects a substantial number of young people. Third, among non-English home language girls, physician services related to depression and anxiety increased substantially (22.1%). Fourth, among English home language boys, physician services related to conduct disorders increased substantially (19.1%). Finally, physician services related to substance use disorders fell

¹² We do not report results for physician services associated with diagnoses in our “other” category because the wide range of conditions in this category make results difficult to interpret. We do not report results for physician services related to eating disorders among boys because of its extremely low pre-COVID baseline in this group.

substantially among English home language boys (-31.1%), Indigenous boys (-43.3%) and Indigenous girls (-27.7%).

We next consider two additional questions: (1) in cases where physician services or prescriptions increased, had treatment levels begun to return to predicted levels by March 2022; and (2) did our before/after estimator miss any cases where an increase began sufficiently late in the period of observation that it did not fully offset the decline in the early months of the pandemic? To investigate these questions, we estimate an event-study model:

$$y_{it} = \sum_{k=0}^K \beta_k \times 1(e_t = k) + \sum_{k=-7}^{-2} \beta_k \times 1(e_t = k) + \beta_{-8} \times 1(e_t \leq -8) \quad (2) \\ + \mu_{m(t)} + \lambda t + \alpha X_{it} + \epsilon_{it}$$

where $e_t = K$ corresponds to March 2022. The parameter β_6 , for example, measures the deviation of the observed share of students who were treated in September 2020 ($e_t = 6$) from its predicted value, relative to the corresponding share in February 2020 ($e_t = -1$). While the magnitudes of event study coefficients are sensitive to the choice of the omitted period, the dynamic patterns they reveal are not. They can therefore be used to determine whether the average post-COVID changes reported in Table 3 persist through March 2022, to identify the timing of any peaks and to determine if there are any cases where increased service use late in the sample period was obscured by the before/after estimator.

A full set of event study plots for all groups and diagnostic categories is available in Appendix Figure B1. Figure 3 presents event study plots for a selected set of cases in which Tables 3a and 3b report a clear (statistically significant at 1%) change in average monthly outcomes. Physician services related to ADHD among Indigenous girls and physician services related to ADHD, depression, psychotic disorders and conduct disorders and prescriptions related to ADHD, depression and psychosis among English home language girls all remained at or near their peak levels in early 2022. The dramatic increase in the frequency of physician visits related to eating disorders among English home language girls appears to have peaked in mid-2021 but remained substantially higher than predicted through March 2022. Physician visits related to depression among other non-English language girls also declined from their peak but remained elevated. The declines in physician services related to depression among English home language boys take the form of a return to predicted levels following a large early decline, while the declines in physician services related to substance abuse for both Indigenous and English home language boys are substantially more persistent.


Figure 3. COVID event-study effects, selected cases.



Each point reports the coefficient on an indicator for a specific month as described in equation (2); control variables include time trend and indicators for month, grade, neighborhood SES quartile and birth quarter. Capped lines indicate 95% confidence intervals. Estimation sample includes observations from September 2016 through March 2022. Cases reported are those where the main effect of COVID reported in Table 3a or Table 3b is statistically significant at 1%.

Inspection of event study plots for the cases where the before-after estimates reported in Table 3 were *not* statistically significant produced one case of interest. As shown in Figure 4, prescriptions related to ADHD among Indigenous girls began to increase substantially relative to predicted rates by late winter of 2020 and continued above predicted levels through to the end of our period of observation.

Figure 4. COVID event-study effects, ADHD-related prescriptions filled, Indigenous girls.



Each point reports the coefficient on an indicator for a specific month as described in equation (2); control variables include time trend and indicators for month, grade, neighborhood SES quartile and birth quarter. Capped lines indicate 95% confidence intervals. Estimation sample includes observations from September 2016 through March 2022. The reported case reported is the only one among girls where the main effect of COVID reported in Table 3 was negative or positive and statistically insignificant and where the pattern of event study coefficients indicates a clear increase in outcomes during the later months of the period of observation period.

We next investigate the role of SES in the effect of COVID on mental health outcomes by adding interactions between the post-COVID indicator and indicators of the quartile of the distribution (across students) of neighborhood SES to the baseline before/after model. A full set of results for the interacted model are presented in Appendix Table B3. Our goal is to investigate whether, in the cases selected for reporting in Figure 3, this effect contributed to narrowing or widening any pre-pandemic socioeconomic gaps in outcomes. The results in Tables 4a and 4b suggest that, in most cases, the changes in physician services that we observe were experienced by students throughout the SES distribution. To the extent that these effects differ across SES quartiles, they tend to be greater among higher SES students, notably in the case of eating disorders among English home language girls. The results in Table 4c show a very different pattern – the increases in prescriptions among English home language girls were not experienced by girls in the top half of the SES distribution, reinforcing the tendency for lower SES girls to fill more prescriptions for these medications.¹³

¹³ The tendency for doctors to disproportionately prescribe anti-psychotic medications to low- versus higher-income children and adolescents has been documented and investigated by Currie et al. (2022) in the Canadian province of Ontario during the pre-pandemic period.

Table 4a: COVID effect on physician services by SES quartile, selected cases (girls).

COVID effect on monthly share receiving physician services (as % of pre-COVID mean)							
Description	Indigenous girls	Non-Indigenous, English home language girls					Non-Indigenous, non-English home language girls
	ADHD	Depression/ Anxiety	ADHD	Psychosis	Conduct	Eating	Depression/ Anxiety
COVID effect by SES quartile							
Q1 (bottom)	24.6*** (8.5)	12.1*** (3.2)	21.0** (8.4)	23.7** (9.2)	112.7*** (28.6)	20.7 (14.7)	10.8 (8.3)
Q2	33.6** (14.0)	14.4*** (3.1)	25.8*** (8.1)	22.6*** (8.4)	39.7** (20.6)	42.9*** (16.3)	19.0** (7.7)
Q3	48.8*** (17.2)	10.1*** (2.9)	15.6** (7.7)	7.8 (7.5)	36.5** (15.9)	51.6*** (15.2)	16.2** (8.2)
Q4	52.2** (25.9)	15.5*** (2.9)	48.0*** (8.6)	17.3** (7.2)	41.3** (17.5)	71.7*** (16.8)	37.4*** (8.3)
P-value for constant effect	0.475	0.506	0.023	0.374	0.111	0.117	0.097
Baseline coefficients on SES quartile indicators							
Q2	40.9** (9.3)	-8.1*** (2.5)	-8.1 (6.3)	-21.4*** (7.5)	-33.3** (15.9)	-0.5 (10.9)	9.2 (5.8)
Q3	51.6*** (11.6)	-9.3*** (2.6)	-10.4 (6.6)	-41.2*** (7.2)	-46.0*** (15.9)	10.3 (11.4)	18.9*** (6.4)
Q4	36.0*** (14.2)	-16.2*** (2.6)	2.0 (6.8)	-47.9*** (7.5)	-57.1*** (15.9)	40.8*** (12.0)	30.9*** (6.8)

Table reports coefficients on the interactions of the post-COVID indicator with the SES quartile indicators (upper panel) and coefficients on SES quartile indicators (lower panel); other control variables include time trend and indicators for month, grade and birth quarter. "P-value for constant effect" refers to a test of the restriction that the four interaction terms are equal. Estimation sample includes observations from September 2016 through March 2022, with sample sizes as reported in Table 3a. Cases reported are those where the main effect of COVID reported in Table 3a is statistically significant at 1%. Standard errors clustered by student in parentheses, *** = 1%, ** = 5%, * = 10%.

Table 4b: COVID effect on physician services by SES quartile, selected cases (boys).

Description	COVID effect on monthly share receiving physician services (as % of pre-COVID mean)			
	Indigenous boys	Non-Indigenous, English home language boys		
	Substance	Depression/ Anxiety	Conduct	Substance
COVID effect by SES quartile				
Q1 (bottom)	-54.3*** (14.9)	-3.6 (4.0)	32.4** (13.2)	-37.7** (16.4)
Q2	-16.3 (21.2)	-7.9** (4.0)	31.6** (13.2)	-24.6 (21.3)
Q3	-56.7** (26.0)	-6.0 (3.8)	11.0 (11.0)	-36.1** (16.4)
Q4	-15.9 (43.8)	-7.5* (4.0)	5.9 (8.8)	-26.2 (16.4)
P-value for constant effect	0.251	0.825	0.145	0.911
Baseline coefficients on SES quartile indicators				
Q2	-8.2 (12.0)	0.3 (3.3)	-9.6 (11.8)	-1.6 (13.1)
Q3	-6.3 (22.1)	-0.1 (3.4)	-14.7 (11.8)	-21.3* (13.1)
Q4	12.0 (32.7)	10.9*** (3.6)	-59.6*** (10.3)	-32.8** (13.1)

Table reports coefficients on the interactions of the post-COVID indicator with the SES quartile indicators (upper panel) and coefficients on SES quartile indicators (lower panel); other control variables include time trend and indicators for month, grade and birth quarter. “P-value for constant effect” refers to a test of the restriction that the four interaction terms are equal. Estimation sample includes observations from September 2016 through March 2022, with sample sizes as reported in Table 3a. Cases reported are those where the main effect of COVID reported in Table 3a is statistically significant at 1%. Standard errors clustered by student in parentheses, *** = 1%, ** = 5%, * = 10%.

Table 4c: COVID effect on prescriptions filled by SES quartile, selected cases.

Description	COVID effect on monthly share filling prescriptions (as % of pre-COVID mean)		
	Non-Indigenous, English home language girls		Non-Indigenous, English home language boys
	Antidepressant and related	Antipsychotic	ADHD-related
COVID effect by SES quartile			
Q1(bottom)	8.6** (3.8)	52.5*** (11.5)	-3.4 (4.6)
Q2	10.0*** (3.5)	21.6** (9.4)	-7.9* (4.1)
Q3	3.3 (3.2)	4.2 (8.6)	-8.3** (3.7)
Q4	0.2 (3.1)	2.9 (7.5)	-13.6*** (3.6)
P-value for constant effect	0.101	0.001	0.335
Baseline coefficients on SES quartile indicators			
Q2	-19.3*** (3.2)	-25.1*** (9.2)	-9.9** (4.6)
Q3	-25.8*** (3.3)	-30.3*** (9.6)	-20.1*** (4.6)
Q4	-36.4*** (3.3)	-51.5*** (9.0)	-26.3*** (4.5)

Table reports coefficients on the interactions of the post-COVID indicator with the SES quartile indicators (upper panel) and coefficients on SES quartile indicators (lower panel); other control variables include time trend and indicators for month, grade and birth quarter. “P-value for constant effect” refers to a test of the restriction that the four interaction terms are equal. Estimation sample includes observations from September 2016 through March 2022, with sample sizes as reported in Table 3b. Cases reported are those where the main effect of COVID reported in Table 3b is statistically significant at 1%. Standard errors clustered by student in parentheses, *** = 1%, ** = 5%, * = 10%.

4.2 In-person versus hybrid learning

Following a province-wide school shut-down in March 2020, the Ministry of Education tasked school districts with developing learning plans for the 2020/21 school year that would keep schools open while incorporating measures to reduce the duration and frequency of contact among students. Districts were permitted to deliver up to 50% of each student’s instructional time in a remote (online) format, although this approach was discouraged for younger grades. This policy resulted in a patchwork of hybrid programs as some districts offered in-person learning only, districts that offered hybrid learning did so in different grades and districts that offered hybrid learning did so for different shares of the school week.

Hybrid learning was most commonly adopted by larger school districts serving urban and suburban communities.¹⁴ In light of evidence that urban and rural communities had very different experiences during COVID (e.g., Cabot and Bushnik 2022), we focus our investigation of the potential mediating role of hybrid learning on variation in the effect of COVID across grades rather than across districts.¹⁵ To isolate this effect, we create two measures of hybrid learning: (i) a (time-invariant) indicator that a district introduced hybrid learning in any grade; and (ii) a (time-invariant) indicator that a student's district introduced hybrid learning to the student's own grade. We include these measures and their interactions with the post-COVID indicator, along with interactions between grade-specific indicators and the post-COVID indicator, in our before/after model. The coefficient on the interaction between the post-COVID indicator and the second hybrid learning indicator (i.e., that a student's district introduced hybrid learning to the student's own grade) captures differences in the response to COVID among students who were exposed to hybrid learning and students in the same set of districts who were not (because hybrid learning was not offered to their grade), while allowing for differences in the response to COVID across grades that is common to all districts. Note that the validity of this estimator rests on the assumption that the underlying trend and seasonal patterns of each outcome variable are the same across grades and hybrid versus non-hybrid school districts.

For each of the selected cases reported in Table 3 (i.e., where we have found clear evidence of a change in mental health services or prescriptions in a specific population group), Table 5 shows the estimates for the post-COVID indicator and its various interactions. The final row of each table shows the coefficient of interest. The effect is statistically significant in only one of the fourteen cases - conduct disorders for English home language boys - and in most cases the point estimates are small. These results provide no evidence that hybrid learning played a meaningful mediating role in the increases in mental health services among girls reported in Table 3. A full set of results for the interacted model are presented in Appendix Table B4.

¹⁴ For example, while only 21 of the 60 school districts adopted hybrid learning for grade 10 students, these districts include all of the largest 11 districts and account for over 70% of all grade 10 students. See Appendix A for further details.

¹⁵ Given that some students enroll in private school, distance learning programs and home-schooling, and some students are enrolled in a grade other than the one indicated by their date of birth (due to, for example, red-shirting), the hybrid learning variables are correlated with but not identical to the actual learning environments of each student.

Table 5a: COVID effects on physician services by learning mode, selected cases (girls).

Description	COVID effect on monthly share receiving physician services (as % of pre-COVID mean)						
	Indigenous girls	Non-Indigenous, English home language girls					Non-Indigenous, non-English home language girls
	ADHD	Depression/ Anxiety	ADHD	Psychosis	Conduct	Eating	Depression/ Anxiety
COVID effect w/interaction terms							
Post x (grade 8)	26.7** (13.6)	6.5* (3.4)	2.0 (10.0)	24.2** (9.5)	107.9*** (28.6)	7.1 (16.3)	31.9* (18.8)
Post x (grade 9)	7.5 (11.7)	13.1*** (3.5)	18.1** (9.0)	22.8** (9.7)	71.4*** (25.4)	42.9** (17.9)	44.7** (19.3)
Post x (grade 10)	18.5 (11.6)	23.4*** (3.7)	38.0*** (9.5)	15.6 (10.3)	52.4** (25.4)	60.9*** (20.1)	50.2** (20.8)
Post x (grade 11)	17.7 (11.6)	21.3*** (3.7)	40.0*** (9.5)	18.9* (10.0)	38.1 (25.4)	26.1 (20.1)	73.1*** (20.3)
Post x (hybrid in any grade)	40.3* (22.0)	-1.7 (3.7)	11.5 (11.1)	-6.4 (9.7)	-34.9 (28.6)	20.7 (18.5)	-33.6* (19.2)
Post x (hybrid in own grade)	-8.7 (23.4)	-3.8 (3.8)	-9.0 (11.5)	3.1 (10.0)	25.4 (30.2)	0.0 (20.1)	6.2 (10.3)

Table reports coefficient on interactions of post-COVID indicator with grade and learning mode indicators; control variables include time trend and indicators for month, grade, neighborhood SES quartile, birth quarter, and any-grade and own-grade hybrid indicators. Estimation sample includes observations from September 2016 through March 2022, with sample sizes as reported in Table 3a. Cases reported are those where the main effect of COVID on physician services reported in Table 3a is statistically significant at 1%. Standard errors clustered by student in parentheses. *** = 1%, ** = 5%, * = 10%.

Table 5b: COVID effects on physician services by learning mode, selected cases (boys).

Description	COVID effect on monthly share receiving physician services (as % of pre-COVID mean)			
	Indigenous boys	Non-Indigenous, English home language boys		
	Substance	Depression/ Anxiety	Conduct	Substance
COVID effect w/interaction terms				
Post x (grade 8)	-31.7** (14.4)	-3.6 (4.4)	33.1** (15.4)	-41.0*** (14.8)
Post x (grade 9)	-46.6** (18.3)	-5.4 (4.3)	34.6** (14.7)	-44.3** (16.4)
Post x (grade 10)	-63.5*** (21.6)	0.6 (4.7)	-13.2 (13.2)	-45.9** (19.7)
Post x (grade 11)	-63.5*** (22.1)	7.9 (4.9)	-0.7 (12.5)	-49.2** (24.6)
Post x (hybrid in any grade)	10.6 (20.2)	-4.9 (4.7)	-38.2** (15.4)	16.4 (13.1)
Post x (hybrid in own grade)	12.5 (24.0)	-5.9 (5.1)	69.9*** (16.2)	4.9 (16.4)

Table reports coefficient on interactions of post-COVID indicator with grade and learning mode indicators; control variables include time trend and indicators for month, grade, neighborhood SES quartile, birth quarter, and any-grade and own-grade hybrid indicators. Estimation sample includes observations from September 2016 through March 2022, with sample sizes as

reported in Table 3a. Cases reported are those where the main effect of COVID on physician services reported in Table 3a is statistically significant at 1%. Standard errors clustered by student in parentheses. *** = 1%, ** = 5%, * = 10%.

Table 5c: COVID effects on prescriptions filled by learning mode, selected cases.

Description	COVID effect on monthly share filling prescriptions (as % of pre-COVID mean)		
	Non-Indigenous, English home language girls		Non-Indigenous, English home language boys
	Antidepressant and related	Antipsychotic	ADHD-related
COVID effect w/interaction terms			
Post x (grade 8)	3.4 (3.8)	40.2*** (10.3)	1.5 (5.6)
Post x (grade 9)	12.1*** (3.9)	54.0*** (10.9)	-2.8 (4.8)
Post x (grade 10)	23.0*** (4.2)	51.0*** (12.3)	-7.5* (4.3)
Post x (grade 11)	26.9*** (4.2)	45.8*** (11.9)	-2.8 (4.3)
Post x (hybrid in any grade)	-14.0*** (4.1)	-33.5*** (11.5)	-9.2 (6.2)
Post x (hybrid in own grade)	-2.7 (4.2)	-12.6 (11.1)	1.6 (6.1)

Table reports coefficient on interaction of post-COVID indicator with grade and learning mode indicators; control variables include time trend and indicators for month, grade, neighborhood SES quartile, birth quarter, and any-grade and own-grade hybrid indicators. Estimation sample includes observations from September 2016 through March 2022, with sample sizes as reported in Table 3b. Cases reported are those where the main effect of COVID on prescription medications reported in Table 3b is statistically significant at 1%. Standard errors clustered by student in parentheses. *** = 1%, ** = 5%, * = 10%.

5 Conclusion

The results in this paper suggest that the mental health of adolescent girls in British Columbia suffered significantly through the first two years of the pandemic. This finding is consistent with similar studies that have found increased use of mental health services in diverse jurisdictions that include Korea (Kim et al. 2022), Norway (Evensen et al. 2022), Ontario (Saunders et al. 2022) and Australia (Costa 2022).

Increases in physician services related to depression and anxiety, the most common diagnosis among adolescent girls in the four years preceding the pandemic, were pronounced and widespread. Apart from the obvious suffering that this increase reflects, adolescent depression has been causally linked to a variety of long-term consequences, including educational attainment, employment, earnings and criminal activity (e.g., Fletcher 2010, 2013; Goodman et al. 2011; Lundborg et al. 2014; Anderson et al. 2015).

These short-run effects of depression on outcomes may understate the overall long-run effects of pandemic-related depression on girls - having a greater share of same-gender peers with depression has been found to increase own rates of depression among girls (Fruehwirth et al. 2019; Giulietti et al. 2022)

and higher rates of depression, lower college attendance and a lower likelihood of working and lower income among adult females (Giuliettei et al. 2022).

Likewise, ADHD has been associated with poor long-term outcomes including lower educational attainment and greater likelihood of welfare receipt and criminal activity (e.g., Currie and Stabile 2006; Fletcher and Wolfe 2008, 2009; Currie et al. 2010; Fletcher 2014). The increased use of physician services and prescription medications for ADHD suggests that adolescent girls with ADHD experienced difficulties during the pandemic. These girls may face a longer road than others to recovery from the learning and social losses experienced during COVID. At the same time, however, the increase in physician services during the pandemic may reflect increased recognition of dysfunction among girls who were previously under-diagnosed. Underdiagnosis of ADHD is thought to be particularly common among girls because they are more likely than boys to experience comorbid depression that masks their ADHD symptoms (Quinn and Madhoo 2014), because physicians may use different criteria when diagnosing ADHD in boys versus girls (e.g., Bruchmuller et al. 2012; Hinshaw 2018; Marquardt 2022) and because girls with ADHD tend to be inattentive rather than impulsive or hyperactive and therefore are less likely than boys to be disruptive in class.¹⁶ To the extent that the pandemic brought this disorder to light for some girls, the associated access to appropriate supports and treatment may serve as a silver lining.

Eating disorders are potentially life-threatening conditions that affect a person's emotional and physical health in a severe way. Disordered patterns of eating tend to be persistent and may lead to hospitalization and death (Ham et al. 2013). The marked increase in the use of physician services in relation to eating disorders among high-SES English home language girls during the first two years of the pandemic is therefore of significant concern despite the relatively small numbers affected. Again, however, some of this increase in physician services may reflect previously undiagnosed cases that came to light as individuals' symptoms deteriorated during the pandemic. Underdiagnosis in general can be an issue in relation to some eating disorders, particularly bulimia nervosa (Ham et al. 2015; Arduini et al. 2019), and earlier diagnosis of eating disorders has been shown to reduce the risk of hospitalization and death. In this sense, again, there may be a silver lining to some cases. At the same time, the relatively low and unchanging frequency of physician services related to eating disorders among non-English home language and Indigenous girls should be interpreted with caution. Evidence from the U.S. shows that, while bulimia is more prevalent among girls who are black or low SES compared to white and high SES girls, these same groups are less likely to have a corresponding diagnosis (Ham et al. 2015). To the extent

¹⁶ Furzer et al. (2022) show that, relative to parents' assessments of children's behavior, the tendency of teachers to overidentify behavior problems among children who are relatively young in grade and under-identify them among children who are relatively old in grade is more pronounced among girls than boys, so that older girls with ADHD are particularly at risk of being missed in the referral and diagnostic process.

that the large gap between treatment for eating disorders among English home language girls and non-English home language and Indigenous girls reflects unmet needs, this gap appears to have grown during the first two years of the pandemic.

The decline in physician visits related to substance use disorders that we find among some groups stands in contrast to the moderate *increase* in visits and hospitalizations related to substance use disorders found by Saunders et al. (2021) in Ontario through February 2021. This difference in results warrants further investigation, particularly in light of the substantially longer periods of school closures in Ontario compared to B.C. (Gallagher-MacKay et al. 2021).

References

- Abe-Kim, Jennifer, David T. Takeuchi, Seunghye Hong, Nolan Zane, Stanley Sue, Michael S. Spencer, Hoa Appel, Ethel Nicdao and Margarita Alegría (2007). Use of mental health-related services among immigrant and US-born Asian Americans: Results from the National Latino and Asian American Study. *American Journal of Public Health* 97(1):91-98.
- Anderson, D. Mark, Resul Cesur and Erdal Tekin (2015). Youth depression and future criminal behavior. *Economic Inquiry* 53(1): 294-317.
- Arduini, Tiziano, Daniela Iorio and Eleonora Patacchini (2019). Weight, reference points, and the onset of eating disorders. *Journal of Health Economics* 65: 170-188.
- Becker, Stephen P., Rosanna Breaux, Caroline N. Cusick, Melissa R. Dvorsky, Nicholas P. Marsh, Emma Sciberras and Joshua M. Langberg (2020). Remote learning during COVID-19: Examining school practices, service continuation, and difficulties for adolescents with and without Attention-Deficit/Hyperactivity Disorder. *Journal of Adolescent Health* 67(6): 769-777.
- Bharadwaj, Prashant, Mallesh M. Pai and Agne Suziedelyte (2017). Mental health stigma. *Economics Letters* 159: 57-60.
- Bruchmuller, Katrin, Jurgen Margraf and Silvia Schneider (2012). Is ADHD diagnosed in accord with diagnostic criteria? Overdiagnosis and influence of client gender on diagnosis. *Journal of Consulting and Clinical Psychology* 80(1): 128-138.
- Busch, Susan H., Ezra Golberstein and Ellen Meara (2014). The FDA And ABCs: Unintended consequences of antidepressant warnings on human capital. *Journal of Human Resources* 49(3):540-571.
- Cabot, Jonathan and Tracey Bushnik (2022). Compliance with precautions to reduce the spread of COVID-19 in Canada. *Health Reports* 33(9): 3-10.

- Cook, Benjamin Lê, Nhi-Ha Trinh, Zhihui Li, Sherry Shu-Yeu Hou and Ana M. Progovac (2017). Trends in racial-ethnic disparities in access to mental health care, 2004–2012. *Psychiatric Services* 68(1): 9-16.
- Costa, Juliana de Oliveira, Malcolm B. Gillies, Andrea L. Schaffer, David Peiris, Helga Zoega and Sallie-Anne Pearson (2022). Changes in antidepressant use in Australia: A nationwide analysis (2015-2021). *Australian and New Zealand Journal of Psychiatry* 57(1):49-57.
- Crawford, Claire, Lorraine Dearden and Ellen Greaves (2011). *Does when you are born matter? The impact of month of birth on children's cognitive and non-cognitive skills in England*. London: Institute for Fiscal Studies. Available at: <https://ifs.org.uk/publications/does-when-you-are-born-matter-impact-month-birth-childrens-cognitive-and-non-cognitive> (accessed: 8 April 2023).
- Currie, Janet and Mark Stabile (2006). Child mental health and human capital accumulation: The case of ADHD. *Journal of Health Economics* 25(6): 1094-1118.
- Currie, Janet and Mark Stabile (2009). Mental health in childhood and human capital. In Jonathan Gruber, ed., *An Economic Perspective on the Problems of Disadvantaged Youth*. Chicago: University of Chicago Press.
- Currie, Janet, Mark Stabile, Phongsack Manivong and Leslie L. Roos (2010). Child health and young adult outcomes. *Journal of Human Resources* 45(3): 517-548.
- Currie, Janet, Paul Kurdyak and Jonathan Zhang (2022). Socioeconomic status and access to mental health care: The case of psychiatric medications for children in Ontario Canada. NBER Working Paper No. 30595. National Bureau of Economic Research.
- Evans, William N., Melinda S. Morrill and Stephen T. Parente (2010). Measuring inappropriate medical diagnosis and treatment in survey data: The case of ADHD among school-age children. *Journal of Health Economics* 29(5): 657-673.
- Evensen, Miriam, Rannveig K. Hart, Anna A. Godøy, Lars J. Hauge, Ingunn O. Lund, Ann K. S. Knudsen, Maja W. Grøtting, Pål Surén and Anne Reneflot (2022). Impact of the COVID-19 pandemic on mental healthcare consultations among children and adolescents in Norway: a nationwide registry study. *European Journal of Child and Adolescent Psychiatry* 32(6):1025-1035
- Fletcher, Jason M. (2010). Adolescent depression and educational attainment: results using sibling fixed effects. *Health Economics* 19(7): 855-871.
- Fletcher, Jason M. (2013). Adolescent depression and adult labor market outcomes. *Southern Economic Journal* 80(1): 26-49.

Fletcher, Jason M. (2014). The effects of childhood ADHD on adult labor market outcomes. *Health Economics* 23(2): 159-181.

Fletcher, Jason and Barbara Wolfe (2008). Child mental health and human capital accumulation: the case of ADHD revisited. *Journal of Health Economics* 27(3): 794-800.

Fletcher, Jason and Barbara Wolfe (2009). Long-term consequences of childhood ADHD on criminal activities. *The Journal of Mental Health Policy and Economics* 12(3): 119-138.

Fruehwirth, Jane Cooley, Sriya Iyer and Anwen Zhang (2019). Religion and depression in adolescence. *Journal of Political Economy* 127(3): 1178-1209.

Fumarco, Luca and Stijn Baert (2019). Relative age effect on European adolescents' social network. *Journal of Economic Behavior & Organization* 168: 318-337.

Furzer, Jill, Elizabeth Dhuey and Audrey Laporte (2022). ADHD misdiagnosis: Causes and mitigators. *Health Economics* 31(9): 1926-1953.

Gadermann, Anne M., Monique Gagné Petteni, Magdalena Janus, Joseph H. Puyat, Martin Guhn and Katholiki Georgiades (2022). Prevalence of mental health disorders among immigrant, refugee, and nonimmigrant children and youth in British Columbia, Canada. *JAMA Network Open* 5(2): e2144934.

Gallagher-MacKay, Kelly, Prachi Srivastava, Kathry Underwood, Elizabeth Dhuey, Lance McCready, Karen B. Born, Antonina Maltsev, Anna Perkhun, Robert Steiner, Kali Barrett and Beate Sander (2021). COVID-19 and education disruption in Ontario: emerging evidence on impacts. *Science Briefs of the Ontario COVID-19 Science Advisory Table* 2(34).

Georgiades, Katholiki, Diana Paksarian, Kara E. Rudolph and Kathleen R. Merikangas (2018). Prevalence of mental disorder and service use by immigrant generation and race/ethnicity among U.S. adolescents. *Journal of the American Academy of Child and Adolescent Psychiatry* 57(4):280-287.

Giulietti, Corrado, Michael Vlassopoulos and Yves Zenou (2022). Peers, gender, and long-term depression. *European Economic Review* 144: 104084.

Goodman, Alissa, Robert Joyce and James P. Smith (2011). The long shadow cast by childhood physical and mental problems on adult life. *Proceedings of the National Academy of Sciences of the USA* 108(15): 6032-6037.

Government of British Columbia (2022a). *Medical Services Plan*.
<https://www2.gov.bc.ca/gov/content/health/health-drug-coverage/msp>

- Government of British Columbia (2022b). *About Pharmanet*.
<https://www2.gov.bc.ca/gov/content/health/health-drug-coverage/pharmacare-for-bc-residents/pharmanet>
- Ham, John C., Daniela Iorio and Michelle Sovinsky (2013). Caught in the bulimic trap? Persistence and state dependence of bulimia among young women. *Journal of Human Resources* 48(3): 736-767.
- Ham, John C., Daniela Iorio and Michelle Sovinsky (2015). Disparities in Bulimia Nervosa: Who is left behind? *Economics Letters* 136: 147–150.
- Hawrilenko, Matt, Emily Kroshus, Pooja Tandon, and Dimitri Christakis (2021). The association between school closures and child mental health during COVID-19. *JAMA network open* 4(9): e2124092-e2124092.
- Hinshaw, Stephen P. (2018). Attention deficit hyperactivity disorder (ADHD): controversy, developmental mechanisms and multiple levels of analysis. *Annual Review of Clinical Psychology* 14: 291-316.
- Hvide, Hans K. and Julian Johnsen (2022). COVID-19 and mental health: a longitudinal population study from Norway. *European Journal of Epidemiology* 37: 167–172.
- Kessler, Ronald C., G. Paul Amminger, Sergio Aguilar-Gaxiola, Jordi Alonso, Sing Lee and T. Bedirhan Üstün (2007). Age of onset of mental disorders: a review of recent literature. *Current Opinion in Psychiatry* 20(4): 359–364.
- Kessler, Ronald C., Shelli Avenevoli and Kathleen Ries Merikangas (2001). Mood disorders in children and adolescents: An epidemiologic perspective. *Biological Psychiatry* 49(12): 1002–1014.
- Kim, So Young, Na-Eun Lee, Dae Myoung Yoo, Ji Hee Kim, Mi Jung Kwon, Joo-Hee Kim, Woo Jin Bang and Hyo Geun Choi (2022). Changes in the mean of medical visits due to psychiatric disease in Korean children and adolescents before and during the COVID-19 pandemic. *Life* 12(4): 600.
- Kim-Cohen, Julia, Avshalom Caspi, Terrie E. Moffitt, HonaLee Harrington, Barry J. Milne and Richie Poulton (2003). Prior juvenile diagnoses in adults with mental disorder: Developmental follow-back of a prospective-longitudinal cohort. *Archives of General Psychiatry* 60(7): 709–717.
- Leong, Christine, Laurence Y. Katz, James M. Bolton et al. (2022). Psychotropic drug use in children and adolescents before and during the COVID-19 pandemic. *JAMA Pediatrics* 176(3): 318–320.
- Lundborg, Petter, Anton Nilsson and Dan-Olof Rooth (2014). Adolescent health and labor market outcomes. *Journal of Health Economics* 37: 25-40.

Marquardt, Kelli (2022). Mis(sed) diagnosis: Physician decision-making and ADHD. Working Paper 2022-23. Federal Reserve Bank of Chicago.

Patalay, Praveetha, Jay Belsky, Peter Fonagy, Panos Vostanis, Neil Humphrey, Jessica Deighton and Miranda Wolpert (2015). The extent and specificity of relative age effects on mental health and functioning in early adolescence. *Journal of Adolescent Health* 57(5): 475-481.

Prinz, Daniel, Michael Chernew, David Cutler and Austin Frakt (2018). Health and economic activity over the lifecycle: Literature review. NBER Working Paper No. 24865. National Bureau of Economic Research.

Quinn, Patricia O. and Manisha Madhoo (2014). A review of attention-deficit/hyperactivity disorder in women and girls: uncovering this hidden diagnosis. *The Primary Care Companion for CNS Disorders* 16(3): PCC.13r01596.

Ridley, Matthew, Gautam Rao, Frank Schilbach and Vikram Patel (2020). Poverty, depression, and anxiety: Causal evidence and mechanisms. *Science* 370(6522): eaay0214.

Saunders, Natasha R., Paul Kurdyak, Therese A. Stukel, Rachel Strauss, Longdi Fu, Jun Guan, Lisa Fiksenbaum, Eyal Cohen, Astrid Guttman, Simone Vigod, Maria Chiu, Charlotte M. Hepburn, Kimberly Moran, William Gardner, Mario Cappelli, Purnima Sundar and Alene Toulany (2022). Utilization of physician-based mental health care services among children and adolescents before and during the COVID-19 pandemic in Ontario, Canada. *JAMA Pediatrics* 176(4): e216298.

Sibley, Margaret H., Mercedes Ortiz, Larissa M. Gaias, Rosemary Reyes, Mahima Joshi, Dana Alexander and Paulo Graziano (2021). Top problems of adolescents and young adults with ADHD during the COVID-19 pandemic. *Journal of Psychiatric Research* 136: 190-197.

Surén, Pål, Anne B. Skirbekk, Lelia Torgersen, Lasse Bang, Anna Godøy and Rannveig K. Hart (2022). Eating disorder diagnoses in children and adolescents in Norway before vs during the COVID-19 pandemic. *JAMA Network Open* 5(7) e2222079.

Svaleryd, Helena, Evelina Björkegren and Jonas Vlachos (2021). The impact of the COVID-19 school closure on adolescents' use of mental healthcare services in Sweden, Working Paper, <https://doi.org/10.1101/2021.12.12.21267684>. Accessed April 8, 2023.

Thompson, Angus H., Roger H. Barnsley and James Battle (2004). The relative age effect and the development of self-esteem. *Educational Research* 46(3): 313-320.

Thornicroft, Graham, Somnath Chatterji, Sara Evans-Lacko, Michael Gruber, Nancy Sampson, Sergio Aguilar-Gaxiola, Ali Al-Hamzawi et al. (2017). Undertreatment of people with major depressive disorder in 21 countries. *The British Journal of Psychiatry* 210(2): 119-124.

Toulany, Arlene, Paul Kurdyak, Therese A. Stukel, Rachel Strauss, Longdi Fu, Jun Guran, Lisa Fiksenbaum, Eyal Cohen, Astrid Gruttman, Simone Vigor, Maria Chiu, Charlotte Moore Hepburn, Kimberly Moran, William Gardner, Mario Cappelli, Purnima Sunday and Natasha Saunders (2023). Sociodemographic differences in physician-based mental health and virtual care utilization and uptake of virtual care among children and adolescents during the COVID-19 Pandemic in Ontario, Canada: A population-based study. *The Canadian Journal of Psychiatry*, in press. doi:10.1177/07067437231156254

Appendix A: Data background

This appendix provides additional background information on how the data were prepared for analysis.

A1 Classification of mental health-related diagnoses and medications

Table A1 below provides details on how we classify mental health diagnoses in the MSP data. Each service in the MSP data has at least one associated diagnosis coded according to the ICD-9 classification system. Our indicator for receiving mental health-related physician service is based on the first three digits of the ICD-9 code for the primary diagnosis associated with the service record.

Table A1: Classification of mental health diagnoses.

Category	Included ICD-9 codes
Depression/anxiety	300 (neurotic disorders), 309 (adjustment reaction), 311 (depressive disorder), 313 (emotional disturbance), 50B (anxiety/depression).
ADHD	314 (hyperkinetic syndrome).
Psychosis	295 (schizophrenia), 296 (affective psychosis), 297 (paranoia), 298 (other nonorganic psychoses), 299 (psychoses of childhood).
Conduct disorder	312 (conduct disturbance).
Substance disorder	291 (psychosis induced by alcohol), 292 (psychosis induced by drugs), 303 (alcohol dependence), 304 (drug dependence), 305 (non-dependent drug abuse).
Eating disorder	3071 (anorexia), 3075 (other eating disorders).
Other	290 (dementia), 293 (delirium not associated with drugs or alcohol), 294 (persistent mental disorders due to conditions classified elsewhere), 301 (personality disorders), 306 (physiological malfunction due to mental factors), 308 (acute reaction to stress), 316 (psychic factors associated with diseases classified elsewhere). 307 (special syndromes or disorders not elsewhere classified) excluding 3071 and 3075.

Table A2 provides details on how we classify mental health-related prescription medications in the Pharmanet data. Each medication dispensed has a drug identification number (DIN) that can then be mapped to the chemical/generic name of the drug. Table A2 provides the basic drug types included in each of our three broad categories, along with chemical/generic names for some commonly-used drugs of each type.

Table A2: Classification of mental health-related prescription medications.

Category	Included drugs
Antidepressant and related	<i>Antidepressants</i> : fluoxetine, sertraline, escitalopram, trazodone, citalopram, venlafaxine, bupropion, etc. <i>Anxiolytics</i> : lorazepam, clonazepam, etc. <i>Sedatives</i> : clobazam, zopiclone, etc.
ADHD-related	<i>Stimulants</i> : methylphenidate, dextroamphetamine, dextroamphetamine sulfate. <i>Agonists</i> : guanfacine. <i>NPRIs</i> : atomoxetine.
Antipsychotic	<i>Antipsychotics</i> : quetiapine, risperidone, aripiprazole, olanzapine, etc.

A2 Classification of school districts by planned learning environment

The analysis in Section 4.2 is based on a classification of B.C. public school districts by their September 2020 plans for in-person versus remote learning environments in each grade. Classifications were constructed by the authors on the basis of detailed plans posted on each district’s website in August and September 2020, along with any district communications to parents and other stakeholders that were also posted to the district’s website at that time. The full set of documents used for classification is available on request from the authors.

1. Based on the available documentation, each district/grade was assigned an estimate of in-person learning time as a percentage of normal class time. In some cases, district plans were coded as range of plausible values rather than a single estimate. For example, a district might say that “at least half” of the school day would be spent in-person, which would be coded as a range of 50-100% in-person time.
2. Every district/grade whose plans implied a value or range of values other than 100% in-person was coded as providing hybrid learning.

Table A3 below reports the full set of districts and grades, along with their planned value or range of in-person times. Note that these in-person times reflect the published initial plans for public schools; actual in-person learning time may have deviated from plans in response to changes in local conditions, enrollment in private school, or participation in distance education or homeschooling programs. Our analysis in Section 4.2 includes all students – public and private – to ensure the before/after comparisons are made over a stable population of students.

Table A3: Districts and grades with hybrid learning plans in 2020/21 academic year. All other districts and grades were planned to be 100% in-person.

School district	Grade(s)	% in-person
23 Central Okanagan	10-12	75
34 Abbotsford	9-12	60
35 Langley	10	70-100
35 Langley	11-12	70
36 Surrey	10-12	66.7
37 Delta	10-12	50-100
38 Richmond	10-12	50
39 Vancouver	8-12	50
40 New Westminster	9-12	80
41 Burnaby	10-12	70
42 Maple Ridge	10-12	70-100
43 Coquitlam	9-12	70
44 North Vancouver	10-12	50-100
45 West Vancouver	10-12	50-100
54 Bulkley Valley	10-12	50-100
61 Greater Victoria	9-12	70
62 Sooke	9-12	50
63 Saanich	9-12	75-100
67 Okanagan Skaha	10-12	75
68 Nanaimo-Ladysmith	8-12	70-100
75 Mission	7-9	90
75 Mission	10-12	35-100
92 Nisga'a	K-12	50

Appendix B: Additional results

Tables B1a-B1b show monthly pre-COVID shares of students receiving mental health-related physician services or filling mental health-related prescriptions. Tables B2a-B2b reports the coefficient estimates and standard errors for our main results. The results in Tables 3a-3b are calculated by combining the results in these four tables. For example, Table B2a indicates a 0.41 percentage point increase in physician services related to depression/anxiety for English home language girls, with a standard error of 0.06. The corresponding pre-COVID mean reported in Table B1a is 3.14%, so the result reported in Table 3 is that the outcome has increased by $100\% * 0.41/3.14 \approx 13.1\%$, with a standard error of $100\% * 0.06/3.14 \approx 1.8\%$.

Figures B1a-B1f show the full set of event study results. Selected results from these figures are reported in Figure 3 and discussed in Section 4 of the main text.

Tables B3a-B3l show the full set of results by SES quartile. Selected results from these tables are reported in Table 4 and discussed in Section 4 of the main text.

Tables B4a-B4l show the full set of results by learning mode. Selected results from these tables are reported in Table 5 and discussed in Section 4.2 of the main text.

Table B1a: Monthly share of students receiving mental health-related physician services by condition, pre-COVID (2016/17 through 2018/19).

Sub-population	Monthly % receiving physician services						
	Depression/ Anxiety	ADHD	Psychosis	Conduct	Substance	Eating	Other
Girls							
Indigenous	4.37	0.74	0.66	0.16	0.32	0.12	0.41
Non-indigenous by home language							
English	3.14	0.44	0.36	0.06	0.07	0.18	0.28
Non-English	1.10	0.13	0.16	0.02	0.02	0.09	0.15
Boys							
Indigenous	1.79	1.46	0.48	0.25	0.21		0.21
Non-indigenous by home language							
English	1.46	1.11	0.37	0.14	0.06		0.16
Non-English	0.61	0.32	0.18	0.05	0.03		0.10

Population consists of monthly observations of B.C. resident students ages 13-16 on December 31 of the current academic year. Percent receiving mental health services is based on services received during current month. See Appendix A for definitions of diagnostic categories.

Table B1b: Monthly share of students filling mental health-related prescriptions by drug category, pre-COVID (2016/17 through 2018/19).

Sub-population	Monthly % filling prescriptions		
	Antidepressant and related	ADHD-related	Antipsychotic
Girls			
Indigenous	4.96	1.92	1.10
Non-indigenous by home language			
English	3.46	0.92	0.52
Non-English	0.86	0.18	0.13
Boys			
Indigenous	2.43	3.75	1.32
Non-indigenous by home language			
English	1.85	2.33	0.56
Non-English	0.45	0.44	0.17

Population consists of monthly observations of B.C. resident students ages 13-16 on December 31 of the current academic year. Percent filling mental health prescription is based on prescriptions filled during current month. See Appendix A for definitions of drug categories.

Table B2a: COVID effect on mental health-related physician services by sub-population, as percentage point change.

Sub-population	COVID effect on monthly % receiving physician services					
	Depression/ Anxiety	ADHD	Psychosis	Conduct	Substance	Eating
Girls						
Indigenous	0.13 (0.14)	0.24*** (0.05)	0.00 (0.05)	0.03 (0.02)	-0.09** (0.04)	0.05 (0.03)
Non-indigenous by home language						
English	0.41*** (0.06)	0.13*** (0.02)	0.06*** (0.02)	0.04*** (0.01)	0.00 (0.01)	0.09*** (0.02)
Non-English	0.24*** (0.05)	0.02 (0.02)	0.01 (0.02)	0.00 (0.01)	-0.01 (0.01)	0.02 (0.02)
Boys						
Indigenous	-0.20** (0.09)	0.02 (0.07)	-0.01 (0.05)	0.05 (0.03)	-0.09*** (0.03)	
Non-indigenous by home language						
English	-0.09*** (0.04)	0.00 (0.03)	0.00 (0.02)	0.03*** (0.01)	-0.02*** (0.01)	
Non-English	0.04 (0.04)	0.01 (0.03)	0.01 (0.02)	0.01 (0.01)	0.01 (0.01)	


Table reports coefficient on post-COVID indicator in percentage point terms; control variables include time trend and indicators for month, grade, neighborhood SES quartile, and birth quarter. Estimation sample includes observations from September 2016 through March 2022, with reported sample size based on number of unique student-month pairs and rounded to nearest 100. Eating disorder results are not reported for boys due to limited variation in the dependent variable. Standard errors clustered by student in parentheses. *** = 1%, ** = 5%, * = 10%.

Table B2b: COVID effect on mental health-related prescriptions filled by sub-population, as percentage point change.

Sub-population	COVID effect on monthly % filling prescriptions		
	Antidepressant and related	ADHD-related	Antipsychotic
Girls			
Indigenous	0.02 (0.17)	0.14 (0.10)	0.04 (0.09)
Non-indigenous by home language			
English	0.18*** (0.07)	0.07** (0.03)	0.10*** (0.03)
Non-English	0.02 (0.06)	-0.05** (0.02)	0.03 (0.02)
Boys			
Indigenous	-0.16 (0.12)	-0.19 (0.14)	0.07 (0.09)
Non-indigenous by home language			
English	-0.09** (0.05)	-0.20*** (0.05)	0.01 (0.03)
Non-English	0.03 (0.04)	-0.07** (0.03)	-0.03 (0.02)


Table reports coefficient on post-COVID indicator in percentage point terms; control variables include time trend and indicators for month, grade, neighborhood SES quartile and birth quarter. Estimation sample includes observations from September 2016 through March 2022, with reported sample size based on number of unique student-month pairs and rounded to nearest 100. Standard errors clustered by student in parentheses. *** = 1%, ** = 5%, * = 10%.

Figure B1a: COVID event-study effects, Indigenous girls.




Each point reports the coefficient on an indicator for a specific month as described in equation (2); control variables include time trend and indicators for month, grade, neighborhood SES quartile and birth quarter. Capped lines indicate 95% confidence intervals. Estimation sample includes observations from September 2016 through March 2022.

Figure B1b: COVID event-study effects, non-Indigenous English home language girls.




Each point reports the coefficient on an indicator for a specific month as described in equation (2); control variables include time trend and indicators for month, grade, neighborhood SES quartile and birth quarter. Capped lines indicate 95% confidence intervals. Estimation sample includes observations from September 2016 through March 2022.

Figure B1c: COVID event-study effects, non-Indigenous non-English home language girls.




Each point reports the coefficient on an indicator for a specific month as described in equation (2); control variables include time trend and indicators for month, grade, neighborhood SES quartile and birth quarter. Capped lines indicate 95% confidence intervals. Estimation sample includes observations from September 2016 through March 2022.

Figure B1d: COVID event-study effects, Indigenous boys




Each point reports the coefficient on an indicator for a specific month as described in equation (2); control variables include time trend and indicators for month, grade, neighborhood SES quartile and birth quarter. Capped lines indicate 95% confidence intervals. Estimation sample includes observations from September 2016 through March 2022.

Figure B1e: COVID event-study effects, non-Indigenous English home language boys.



Each point reports the coefficient on an indicator for a specific month as described in equation (2); control variables include time trend and indicators for month, grade, neighborhood SES quartile and birth quarter. Capped lines indicate 95% confidence intervals. Estimation sample includes observations from September 2016 through March 2022.

Figure B1f: COVID event-study effects, non-Indigenous non-English home language boys.



Each point reports the coefficient on an indicator for a specific month as described in equation (2); control variables include time trend and indicators for month, grade, neighborhood SES quartile and birth quarter. Capped lines indicate 95% confidence intervals. Estimation sample includes observations from September 2016 through March 2022.

Table B3a: COVID effects on physician services by SES quartile, Indigenous girls.

Description	COVID effect on monthly % receiving physician services					
	Depression/ Anxiety	ADHD	Psychosis	Conduct	Substance	Eating
COVID effect by SES quartile						
Q1(bottom)	-0.12 (0.16)	0.18*** (0.06)	0.00 (0.06)	0.03 (0.03)	-0.10** (0.04)	0.03 (0.03)
Q2	0.50** (0.24)	0.25** (0.10)	0.00 (0.08)	0.04 (0.04)	-0.09* (0.05)	0.16*** (0.05)
Q3	0.68** (0.30)	0.36*** (0.13)	0.03 (0.10)	0.03 (0.05)	-0.09* (0.05)	0.03 (0.07)
Q4	-0.16 (0.43)	0.39** (0.19)	-0.05 (0.17)	0.04 (0.06)	-0.02 (0.09)	-0.19* (0.11)
P-value for constant effect	0.018	0.475	0.980	0.991	0.850	0.015
Baseline coefficients on SES quartile indicators						
Q2	0.32** (0.16)	0.30*** (0.07)	0.08 (0.06)	0.05* (0.03)	-0.02 (0.03)	0.02 (0.03)
Q3	0.52*** (0.20)	0.38*** (0.09)	-0.02 (0.08)	0.04 (0.03)	-0.08** (0.04)	0.13*** (0.05)
Q4	0.82*** (0.31)	0.27** (0.11)	0.11 (0.14)	0.03 (0.04)	-0.03 (0.07)	0.22** (0.10)

Table reports coefficients on the interactions of the post-COVID indicator with the SES quartile indicators (upper panel) and coefficients on SES quartile indicators (lower panel); other control variables include time trend and indicators for month, grade and birth quarter. "P-value for constant effect" refers to a test of the restriction that the four interaction terms are equal. Estimation sample includes observations from September 2016 through March 2022, with sample sizes as reported in Table 3a. Standard errors clustered by student in parentheses. *** = 1%, ** = 5%, * = 10%.

Table B3b: COVID effects on prescriptions filled by SES quartile, Indigenous girls.

Description	COVID effect on monthly % filling prescriptions		
	Antidepressant and related	ADHD-related	Antipsychotic
COVID effect by SES quartile			
Q1(bottom)	-0.41* (0.21)	0.06 (0.13)	0.04 (0.11)
Q2	0.78** (0.32)	0.11 (0.21)	0.06 (0.17)
Q3	0.69* (0.39)	0.22 (0.28)	0.08 (0.18)
Q4	-0.50 (0.54)	0.64* (0.38)	-0.15 (0.26)
P-value for constant effect	0.001	0.533	0.891
Baseline coefficients on SES quartile indicators			
Q2	0.35 (0.23)	0.50*** (0.18)	0.21 (0.13)
Q3	0.13 (0.28)	0.68*** (0.23)	-0.12 (0.14)
Q4	0.53 (0.41)	0.12 (0.28)	-0.10 (0.20)

Table reports coefficients on the interactions of the post-COVID indicator with the SES quartile indicators (upper panel) and coefficients on SES quartile indicators (lower panel); other control variables include time trend and indicators for month, grade and birth quarter. "P-value for constant effect" refers to a test of the restriction that the four interaction terms are equal. Estimation sample includes observations from September 2016 through March 2022, with sample sizes as reported in Table 3b. Standard errors clustered by student in parentheses. *** = 1%, ** = 5%, * = 10%.

Table B3c: COVID effects on physician services by SES quartile, non-Indigenous English home language girls.

Description	COVID effect on monthly % receiving physician services					
	Depression/ Anxiety	ADHD	Psychosis	Conduct	Substance	Eating
COVID effect by SES quartile						
Q1(bottom)	0.38*** (0.10)	0.09** (0.04)	0.09** (0.03)	0.07*** (0.02)	-0.02 (0.01)	0.04 (0.03)
Q2	0.45*** (0.10)	0.11*** (0.04)	0.08*** (0.03)	0.03** (0.01)	0.00 (0.01)	0.08*** (0.03)
Q3	0.32*** (0.09)	0.07** (0.03)	0.03 (0.03)	0.02** (0.01)	-0.01 (0.01)	0.10*** (0.03)
Q4	0.49*** (0.09)	0.21*** (0.04)	0.06** (0.03)	0.03** (0.01)	0.02** (0.01)	0.13*** (0.03)
P-value for constant effect	0.506	0.023	0.374	0.111	0.026	0.117
Baseline coefficients on SES quartile indicators						
Q2	-0.25*** (0.08)	-0.04 (0.03)	-0.08*** (0.03)	-0.02** (0.01)	-0.02* (0.01)	0.00 (0.02)
Q3	-0.29*** (0.08)	-0.05 (0.03)	-0.15*** (0.03)	-0.03*** (0.01)	-0.04*** (0.01)	0.02 (0.02)
Q4	-0.51*** (0.08)	0.01 (0.03)	-0.17*** (0.03)	-0.04*** (0.01)	-0.05*** (0.01)	0.08*** (0.02)

Table reports coefficients on the interactions of the post-COVID indicator with the SES quartile indicators (upper panel) and coefficients on SES quartile indicators (lower panel); other control variables include time trend and indicators for month, grade and birth quarter. "P-value for constant effect" refers to a test of the restriction that the four interaction terms are equal. Estimation sample includes observations from September 2016 through March 2022, with sample sizes as reported in Table 3a. Standard errors clustered by student in parentheses. *** = 1%, ** = 5%, * = 10%.

Table B3d: COVID effects on prescriptions filled by SES quartile, non-Indigenous English home language girls.

Description	COVID effect on monthly % filling prescriptions			
	Antidepressant and related	ADHD-related	Antipsychotic	
COVID effect by SES quartile				
	Q1(bottom)	0.30** (0.13)	0.16** (0.07)	0.27*** (0.06)
	Q2	0.35*** (0.12)	0.15** (0.07)	0.11** (0.05)
	Q3	0.11 (0.11)	-0.05 (0.06)	0.02 (0.05)
	Q4	0.01 (0.11)	0.05 (0.06)	0.02 (0.04)
P-value for constant effect		0.101	0.056	0.001
Baseline coefficients on SES quartile indicators				
	Q2	-0.67*** (0.11)	-0.21*** (0.07)	-0.13*** (0.05)
	Q3	-0.89*** (0.12)	-0.28*** (0.07)	-0.16*** (0.05)
	Q4	-1.26*** (0.11)	-0.24*** (0.07)	-0.27*** (0.05)

Table reports coefficients on the interactions of the post-COVID indicator with the SES quartile indicators (upper panel) and coefficients on SES quartile indicators (lower panel); other control variables include time trend and indicators for month, grade and birth quarter. "P-value for constant effect" refers to a test of the restriction that the four interaction terms are equal. Estimation sample includes observations from September 2016 through March 2022, with sample sizes as reported in Table 3b. Standard errors clustered by student in parentheses. *** = 1%, ** = 5%, * = 10%.

Table B3e: COVID effects on physician services by SES quartile, non-Indigenous non-English home language girls.

Description	COVID effect on monthly % receiving physician services					
	Depression/ Anxiety	ADHD	Psychosis	Conduct	Substance	Eating
COVID effect by SES quartile						
Q1(bottom)	0.12 (0.09)	-0.02 (0.03)	0.04 (0.04)	0.02 (0.01)	-0.02* (0.01)	-0.04 (0.02)
Q2	0.21** (0.09)	0.00 (0.02)	0.00 (0.03)	-0.01 (0.01)	-0.01 (0.01)	0.01 (0.02)
Q3	0.18** (0.09)	0.04 (0.03)	0.01 (0.03)	0.00 (0.01)	-0.01 (0.01)	0.03 (0.02)
Q4	0.41*** (0.09)	0.03 (0.03)	-0.01 (0.03)	0.01 (0.01)	0.00 (0.01)	0.05 (0.03)
P-value for constant effect	0.097	0.304	0.712	0.241	0.243	0.030
Baseline coefficients on SES quartile indicators						
Q2	0.10 (0.06)	0.04** (0.02)	0.01 (0.03)	0.01 (0.01)	0.00 (0.01)	0.01 (0.02)
Q3	0.21*** (0.07)	0.08*** (0.02)	0.02 (0.03)	0.00 (0.01)	-0.01 (0.01)	0.02 (0.02)
Q4	0.34*** (0.08)	0.14*** (0.02)	0.07** (0.03)	0.00 (0.01)	-0.02*** (0.01)	0.11*** (0.03)

Table reports coefficients on the interactions of the post-COVID indicator with the SES quartile indicators (upper panel) and coefficients on SES quartile indicators (lower panel); other control variables include time trend and indicators for month, grade and birth quarter. "P-value for constant effect" refers to a test of the restriction that the four interaction terms are equal. Estimation sample includes observations from September 2016 through March 2022, with sample sizes as reported in Table 3a. Standard errors clustered by student in parentheses. *** = 1%, ** = 5%, * = 10%.

Table B3f: COVID effects on prescriptions filled by SES quartile, non-Indigenous non-English home language girls.

Description	COVID effect on monthly % filling prescriptions		
	Antidepressant and related	ADHD-related	Antipsychotic
COVID effect by SES quartile			
Q1(bottom)	0.03 (0.11)	-0.03 (0.04)	0.08* (0.05)
Q2	-0.01 (0.09)	-0.03 (0.04)	0.02 (0.04)
Q3	-0.02 (0.09)	-0.09** (0.04)	0.02 (0.04)
Q4	0.08 (0.09)	-0.04 (0.04)	0.03 (0.04)
P-value for constant effect	0.873	0.638	0.719
Baseline coefficients on SES quartile indicators			
Q2	0.03 (0.08)	0.04 (0.04)	-0.01 (0.04)
Q3	0.06 (0.09)	0.11*** (0.04)	-0.02 (0.04)
Q4	0.20** (0.09)	0.15*** (0.05)	0.06 (0.04)

Table reports coefficients on the interactions of the post-COVID indicator with the SES quartile indicators (upper panel) and coefficients on SES quartile indicators (lower panel); other control variables include time trend and indicators for month, grade and birth quarter. "P-value for constant effect" refers to a test of the restriction that the four interaction terms are equal. Estimation sample includes observations from September 2016 through March 2022, with sample sizes as reported in Table 3b. Standard errors clustered by student in parentheses. *** = 1%, ** = 5%, * = 10%.

Table B3g: COVID effects on physician services by SES quartile, Indigenous boys.

Description	COVID effect on monthly % receiving physician services				
	Depression/ Anxiety	ADHD	Psychosis	Conduct	Substance
COVID effect by SES quartile					
Q1(bottom)	-0.31*** (0.10)	-0.14 (0.09)	-0.02 (0.05)	0.05 (0.03)	-0.11*** (0.03)
Q2	-0.17 (0.14)	0.31** (0.13)	-0.02 (0.07)	0.05 (0.05)	-0.03 (0.04)
Q3	0.07 (0.19)	0.17 (0.18)	0.05 (0.10)	-0.01 (0.08)	-0.12** (0.05)
Q4	0.00 (0.30)	0.02 (0.24)	-0.01 (0.14)	0.11* (0.06)	-0.03 (0.09)
P-value for constant effect	0.217	0.012	0.898	0.674	0.251
Baseline coefficients on SES quartile indicators					
Q2	0.34*** (0.10)	0.26*** (0.09)	0.04 (0.05)	0.08** (0.04)	-0.02 (0.03)
Q3	0.42*** (0.12)	0.47*** (0.12)	0.10 (0.07)	0.13** (0.06)	-0.01 (0.05)
Q4	0.83*** (0.22)	0.68*** (0.20)	0.07 (0.11)	-0.05 (0.04)	0.03 (0.07)

Table reports coefficients on the interactions of the post-COVID indicator with the SES quartile indicators (upper panel) and coefficients on SES quartile indicators (lower panel); other control variables include time trend and indicators for month, grade and birth quarter. "P-value for constant effect" refers to a test of the restriction that the four interaction terms are equal. Estimation sample includes observations from September 2016 through March 2022, with sample sizes as reported in Table 3a. Standard errors clustered by student in parentheses. *** = 1%, ** = 5%, * = 10%.

Table B3h: COVID effects on prescriptions filled by SES quartile, Indigenous boys.

Description	COVID effect on monthly % filling prescriptions		
	Antidepressant and related	ADHD-related	Antipsychotic
COVID effect by SES quartile			
Q1(bottom)	-0.35** (0.15)	-0.30 (0.18)	0.04 (0.11)
Q2	-0.20 (0.23)	-0.12 (0.29)	0.19 (0.18)
Q3	0.67** (0.31)	0.26 (0.37)	0.04 (0.23)
Q4	-0.33 (0.42)	-0.39 (0.52)	0.00 (0.29)
P-value for constant effect	0.027	0.554	0.914
Baseline coefficients on SES quartile indicators			
Q2	0.47*** (0.18)	0.73*** (0.25)	0.26* (0.15)
Q3	0.29 (0.22)	0.75** (0.31)	0.23 (0.19)
Q4	0.76** (0.34)	0.93** (0.46)	-0.17 (0.22)

Table reports coefficients on the interactions of the post-COVID indicator with the SES quartile indicators (upper panel) and coefficients on SES quartile indicators (lower panel); other control variables include time trend and indicators for month, grade and birth quarter. "P-value for constant effect" refers to a test of the restriction that the four interaction terms are equal. Estimation sample includes observations from September 2016 through March 2022, with sample sizes as reported in Table 3b. Standard errors clustered by student in parentheses. *** = 1%, ** = 5%, * = 10%.

Table B3i: COVID effects on physician services by SES quartile, non-Indigenous English home language boys.

Description	COVID effect on monthly % receiving physician services				
	Depression/ Anxiety	ADHD	Psychosis	Conduct	Substance
COVID effect by SES quartile					
Q1(bottom)	-0.05 (0.06)	-0.05 (0.05)	0.03 (0.03)	0.04** (0.02)	-0.02** (0.01)
Q2	-0.12** (0.06)	-0.05 (0.05)	-0.03 (0.03)	0.04** (0.02)	-0.02 (0.01)
Q3	-0.09 (0.06)	0.06 (0.05)	-0.02 (0.03)	0.02 (0.02)	-0.02** (0.01)
Q4	-0.11* (0.06)	0.03 (0.05)	0.00 (0.03)	0.01 (0.01)	-0.02 (0.01)
P-value for constant effect	0.825	0.190	0.537	0.145	0.911
Baseline coefficients on SES quartile indicators					
Q2	0.00 (0.05)	0.00 (0.04)	-0.04 (0.03)	-0.01 (0.02)	0.00 (0.01)
Q3	0.00 (0.05)	-0.09* (0.04)	-0.11*** (0.03)	-0.02 (0.02)	-0.01* (0.01)
Q4	0.16*** (0.05)	0.02 (0.05)	-0.13*** (0.03)	-0.08*** (0.01)	-0.02** (0.01)

Table reports coefficients on the interactions of the post-COVID indicator with the SES quartile indicators (upper panel) and coefficients on SES quartile indicators (lower panel); other control variables include time trend and indicators for month, grade and birth quarter. "P-value for constant effect" refers to a test of the restriction that the four interaction terms are equal. Estimation sample includes observations from September 2016 through March 2022, with sample sizes as reported in Table 3a. Standard errors clustered by student in parentheses. *** = 1%, ** = 5%, * = 10%.

Table B3j: COVID effects on prescriptions filled by SES quartile, non-Indigenous English home language boys.

Description	COVID effect on monthly % filling prescriptions		
	Antidepressant and related	ADHD-related	Antipsychotic
COVID effect by SES quartile			
Q1(bottom)	0.02 (0.09)	-0.08 (0.11)	0.04 (0.06)
Q2	-0.06 (0.09)	-0.18* (0.10)	0.11** (0.05)
Q3	-0.14* (0.08)	-0.19** (0.09)	-0.04 (0.05)
Q4	-0.16** (0.08)	-0.32*** (0.08)	-0.05 (0.04)
P-value for constant effect	0.451	0.335	0.069
Baseline coefficients on SES quartile indicators			
Q2	-0.13 (0.09)	-0.23** (0.11)	-0.19*** (0.06)
Q3	-0.32*** (0.09)	-0.47*** (0.11)	-0.23*** (0.06)
Q4	-0.32*** (0.09)	-0.61*** (0.11)	-0.35*** (0.06)

Table reports coefficients on the interactions of the post-COVID indicator with the SES quartile indicators (upper panel) and coefficients on SES quartile indicators (lower panel); other control variables include time trend and indicators for month, grade and birth quarter. "P-value for constant effect" refers to a test of the restriction that the four interaction terms are equal. Estimation sample includes observations from September 2016 through March 2022, with sample sizes as reported in Table 3b. Standard errors clustered by student in parentheses. *** = 1%, ** = 5%, * = 10%.

Table B3k: COVID effects on physician services by SES quartile, non-Indigenous non-English home language boys.

Description	COVID effect on monthly % receiving physician services				
	Depression/ Anxiety	ADHD	Psychosis	Conduct	Substance
COVID effect by SES quartile					
Q1(bottom)	0.03 (0.06)	-0.05 (0.04)	0.04 (0.04)	0.01 (0.02)	0.00 (0.01)
Q2	0.00 (0.05)	0.04 (0.04)	-0.03 (0.03)	-0.01 (0.02)	0.00 (0.01)
Q3	0.02 (0.06)	-0.01 (0.04)	0.02 (0.04)	0.00 (0.01)	-0.01 (0.01)
Q4	0.10 (0.07)	0.03 (0.04)	0.02 (0.03)	0.01 (0.01)	0.02** (0.01)
P-value for constant effect	0.570	0.185	0.401	0.445	0.071
Baseline coefficients on SES quartile indicators					
Q2	0.01 (0.04)	0.01 (0.03)	0.01 (0.03)	-0.01 (0.01)	0.00 (0.01)
Q3	0.13*** (0.05)	0.10*** (0.03)	0.03 (0.03)	-0.01 (0.01)	0.00 (0.01)
Q4	0.27*** (0.05)	0.19*** (0.04)	0.01 (0.03)	-0.02* (0.01)	-0.01*** (0.01)

Table reports coefficients on the interactions of the post-COVID indicator with the SES quartile indicators (upper panel) and coefficients on SES quartile indicators (lower panel); other control variables include time trend and indicators for month, grade and birth quarter. "P-value for constant effect" refers to a test of the restriction that the four interaction terms are equal. Estimation sample includes observations from September 2016 through March 2022, with sample sizes as reported in Table 3a. Standard errors clustered by student in parentheses. *** = 1%, ** = 5%, * = 10%.

Table B3l: COVID effects on prescriptions filled by SES quartile, non-Indigenous non-English home language boys.

Description	COVID effect on monthly % filling prescriptions		
	Antidepressant and related	ADHD-related	Antipsychotic
COVID effect by SES quartile			
Q1(bottom)	0.07 (0.07)	-0.03 (0.07)	0.00 (0.06)
Q2	-0.07 (0.07)	-0.04 (0.06)	-0.06 (0.04)
Q3	-0.03 (0.06)	-0.12** (0.06)	-0.04 (0.04)
Q4	0.16** (0.06)	-0.07 (0.06)	0.00 (0.03)
P-value for constant effect	0.036	0.672	0.590
Baseline coefficients on SES quartile indicators			
Q2	0.10 (0.06)	0.00 (0.06)	-0.06 (0.06)
Q3	0.09 (0.06)	0.13* (0.07)	-0.07 (0.05)
Q4	0.07 (0.06)	0.16** (0.07)	-0.12** (0.05)

Table reports coefficients on the interactions of the post-COVID indicator with the SES quartile indicators (upper panel) and coefficients on SES quartile indicators (lower panel); other control variables include time trend and indicators for month, grade and birth quarter. "P-value for constant effect" refers to a test of the restriction that the four interaction terms are equal. Estimation sample includes observations from September 2016 through March 2022, with sample sizes as reported in Table 3b. Standard errors clustered by student in parentheses. *** = 1%, ** = 5%, * = 10%.

Table B4a: COVID effects on physician services by learning mode, Indigenous girls.

Description	COVID effect on monthly % receiving physician services					
	Depression/ anxiety	ADHD	Psychosis	Conduct	Substance	Eating
COVID effect w/interaction terms						
Post x (grade 8)	-0.02 (0.22)	0.20** (0.10)	0.10 (0.07)	0.00 (0.03)	-0.04 (0.04)	0.02 (0.04)
Post x (grade 9)	0.06 (0.22)	0.06 (0.09)	0.03 (0.08)	-0.01 (0.03)	-0.07 (0.04)	0.05 (0.04)
Post x (grade 10)	0.04 (0.22)	0.14 (0.09)	-0.02 (0.08)	0.04 (0.04)	-0.14** (0.05)	0.14*** (0.05)
Post x (grade 11)	0.26 (0.23)	0.13 (0.09)	-0.02 (0.09)	0.05 (0.04)	-0.10* (0.06)	0.10** (0.05)
Post x (hybrid in any grade)	0.35 (0.31)	0.30* (0.16)	0.09 (0.10)	0.02 (0.04)	0.06 (0.04)	-0.01 (0.05)
Post x (hybrid in own grade)	-0.33 (0.36)	-0.07 (0.17)	-0.21* (0.12)	0.02 (0.06)	-0.09 (0.06)	-0.09 (0.07)

Table reports coefficient on interactions of post-COVID indicator with grade and learning mode indicators; control variables include time trend and indicators for month, grade, neighborhood SES quartile, birth quarter, and any-grade and own-grade hybrid indicators. Estimation sample includes observations from September 2016 through March 2022, with sample sizes as reported in Table 3a. Standard errors clustered by student in parentheses. *** = 1%, ** = 5%, * = 10%.

Table B4b: COVID effects on prescriptions filled by learning mode, Indigenous girls.

Description	COVID effect on monthly % filling prescriptions		
	Antidepressant and related	ADHD-related	Antipsychotic
COVID effect w/interaction terms			
Post x (grade 8)	-0.26 (0.28)	0.21 (0.22)	-0.01 (0.15)
Post x (grade 9)	-0.19 (0.29)	-0.05 (0.19)	-0.06 (0.15)
Post x (grade 10)	0.09 (0.30)	-0.09 (0.17)	0.26* (0.15)
Post x (grade 11)	0.23 (0.31)	0.34* (0.18)	0.36** (0.17)
Post x (hybrid in any grade)	0.08 (0.39)	-0.18 (0.32)	0.00 (0.20)
Post x (hybrid in own grade)	0.09 (0.46)	0.40 (0.37)	-0.32 (0.24)

Table reports coefficient on interactions of post-COVID indicator with grade and learning mode indicators; control variables include time trend and indicators for month, grade, neighborhood SES quartile, birth quarter, and any-grade and own-grade hybrid indicators. Estimation sample includes observations from September 2016 through March 2022, with sample sizes as reported in Table 3a. Standard errors clustered by student in parentheses. *** = 1%, ** = 5%, * = 10%.

Table B4c: COVID effects on physician services by learning mode, non-Indigenous English home language girls.

Description	COVID effect on monthly % receiving physician services					
	Depression/ anxiety	ADHD	Psychosis	Conduct	Substance	Eating
COVID effect w/interaction terms						
Post x (grade 8)	0.21* (0.11)	0.01 (0.04)	0.09** (0.03)	0.07*** (0.02)	0.00 (0.01)	0.01 (0.03)
Post x (grade 9)	0.41*** (0.11)	0.08** (0.04)	0.08** (0.04)	0.05*** (0.02)	0.00 (0.01)	0.08** (0.03)
Post x (grade 10)	0.73*** (0.12)	0.17*** (0.04)	0.06 (0.04)	0.03** (0.02)	-0.02 (0.01)	0.11*** (0.04)
Post x (grade 11)	0.67*** (0.12)	0.18*** (0.04)	0.07* (0.04)	0.02 (0.02)	-0.02 (0.02)	0.05 (0.04)
Post x (hybrid in any grade)	-0.05 (0.12)	0.05 (0.05)	-0.02 (0.04)	-0.02 (0.02)	0.00 (0.01)	0.04 (0.03)
Post x (hybrid in own grade)	-0.12 (0.12)	-0.04 (0.05)	0.01 (0.04)	0.02 (0.02)	0.01 (0.01)	0.00 (0.04)

Table reports coefficient on interactions of post-COVID indicator with grade and learning mode indicators; control variables include time trend and indicators for month, grade, neighborhood SES quartile, birth quarter, and any-grade and own-grade hybrid indicators. Estimation sample includes observations from September 2016 through March 2022, with sample sizes as reported in Table 3a. Standard errors clustered by student in parentheses. *** = 1%, ** = 5%, * = 10%.

Table B4d: COVID effects on prescriptions filled by learning mode, non-Indigenous English home language girls.

Description	COVID effect on monthly % filling prescriptions		
	Antidepressant and related	ADHD-related	Antipsychotic
COVID effect w/interaction terms:			
Post x (grade 8)	0.12 (0.13)	0.06 (0.09)	0.21*** (0.05)
Post x (grade 9)	0.42*** (0.13)	0.16** (0.08)	0.28*** (0.06)
Post x (grade 10)	0.79*** (0.14)	0.28*** (0.08)	0.27*** (0.06)
Post x (grade 11)	0.93*** (0.15)	0.25*** (0.08)	0.24*** (0.06)
Post x (hybrid in any grade)	-0.49*** (0.14)	-0.25*** (0.09)	-0.18*** (0.06)
Post x (hybrid in own grade)	-0.09 (0.15)	0.12 (0.10)	-0.07 (0.06)

Table reports coefficient on interactions of post-COVID indicator with grade and learning mode indicators; control variables include time trend and indicators for month, grade, neighborhood SES quartile, birth quarter, and any-grade and own-grade hybrid indicators. Estimation sample includes observations from September 2016 through March 2022, with sample sizes as reported in Table 3a. Standard errors clustered by student in parentheses. *** = 1%, ** = 5%, * = 10%.

Table B4e: COVID effects on physician services by learning mode, non-Indigenous non-English home language girls.

Description	COVID effect on monthly % receiving physician services					
	Depression/ anxiety	ADHD	Psychosis	Conduct	Substance	Eating
COVID effect w/interaction terms						
Post x (grade 8)	0.35* (0.21)	0.06 (0.06)	0.12** (0.06)	-0.01 (0.02)	-0.02 (0.02)	0.06 (0.08)
Post x (grade 9)	0.49** (0.21)	0.10* (0.06)	0.14** (0.06)	-0.01 (0.02)	-0.01 (0.02)	0.03 (0.09)
Post x (grade 10)	0.55** (0.23)	0.11* (0.06)	0.13* (0.07)	-0.02 (0.02)	-0.02 (0.02)	0.04 (0.09)
Post x (grade 11)	0.81*** (0.22)	0.17*** (0.06)	0.19*** (0.07)	-0.02 (0.02)	0.01 (0.02)	0.08 (0.09)
Post x (hybrid in any grade)	-0.37* (0.21)	-0.10* (0.06)	-0.17*** (0.06)	0.02 (0.02)	0.01 (0.02)	-0.02 (0.09)
Post x (hybrid in own grade)	0.07 (0.11)	0.00 (0.04)	0.04 (0.04)	0.00 (0.01)	-0.01 (0.01)	-0.02 (0.04)

Table reports coefficient on interactions of post-COVID indicator with grade and learning mode indicators; control variables include time trend and indicators for month, grade, neighborhood SES quartile, birth quarter, and any-grade and own-grade hybrid indicators. Estimation sample includes observations from September 2016 through March 2022, with sample sizes as reported in Table 3a. Standard errors clustered by student in parentheses. *** = 1%, ** = 5%, * = 10%.

Table B4f: COVID effects on prescriptions filled by learning mode, non-Indigenous non-English home language girls.

Description	COVID effect on monthly % filling prescriptions		
	Antidepressant and related	ADHD-related	Antipsychotic
COVID effect w/interaction terms			
Post x (grade 8)	0.10 (0.20)	-0.03 (0.08)	-0.01 (0.07)
Post x (grade 9)	0.20 (0.21)	0.02 (0.08)	0.01 (0.07)
Post x (grade 10)	0.22 (0.23)	0.11 (0.09)	0.11 (0.08)
Post x (grade 11)	0.63*** (0.23)	0.15* (0.09)	0.19** (0.09)
Post x (hybrid in any grade)	-0.32 (0.21)	-0.11 (0.08)	-0.03 (0.07)
Post x (hybrid in own grade)	0.05 (0.11)	-0.01 (0.06)	-0.03 (0.06)

Table reports coefficient on interactions of post-COVID indicator with grade and learning mode indicators; control variables include time trend and indicators for month, grade, neighborhood SES quartile, birth quarter, and any-grade and own-grade hybrid indicators. Estimation sample includes observations from September 2016 through March 2022, with sample sizes as reported in Table 3a. Standard errors clustered by student in parentheses. *** = 1%, ** = 5%, * = 10%.

Table B4g: COVID effects on physician services by learning mode, Indigenous boys.

Description	COVID effect on monthly % receiving physician services				
	Depression/ anxiety	ADHD	Psychosis	Conduct	Substance
COVID effect w/interaction terms					
Post x (grade 8)	-0.31** (0.13)	0.05 (0.14)	0.02 (0.07)	0.02 (0.05)	-0.07** (0.03)
Post x (grade 9)	-0.27** (0.13)	0.02 (0.12)	-0.04 (0.07)	0.05 (0.04)	-0.10** (0.04)
Post x (grade 10)	-0.10 (0.14)	-0.09 (0.10)	-0.03 (0.07)	-0.02 (0.04)	-0.13*** (0.05)
Post x (grade 11)	-0.32** (0.14)	-0.36*** (0.10)	-0.07 (0.07)	0.03 (0.04)	-0.13*** (0.05)
Post x (hybrid in any grade)	-0.07 (0.19)	0.44** (0.22)	0.13 (0.11)	0.03 (0.07)	0.02 (0.04)
Post x (hybrid in own grade)	0.29 (0.22)	-0.26 (0.24)	-0.11 (0.13)	0.06 (0.08)	0.03 (0.05)

Table reports coefficient on interactions of post-COVID indicator with grade and learning mode indicators; control variables include time trend and indicators for month, grade, neighborhood SES quartile, birth quarter, and any-grade and own-grade hybrid indicators. Estimation sample includes observations from September 2016 through March 2022, with sample sizes as reported in Table 3a. Standard errors clustered by student in parentheses. *** = 1%, ** = 5%, * = 10%.

Table B4h: COVID effects on prescriptions filled by learning mode, Indigenous boys.

Description	COVID effect on monthly % filling prescriptions		
	Antidepressant and related	ADHD-related	Antipsychotic
COVID effect w/interaction terms			
Post x (grade 8)	-0.43* (0.23)	-0.28 (0.32)	0.11 (0.19)
Post x (grade 9)	-0.30 (0.21)	-0.10 (0.26)	0.25 (0.16)
Post x (grade 10)	-0.10 (0.20)	-0.20 (0.21)	0.31** (0.15)
Post x (grade 11)	-0.48** (0.21)	-0.62*** (0.21)	0.06 (0.15)
Post x (hybrid in any grade)	0.12 (0.34)	0.18 (0.48)	-0.27 (0.26)
Post x (hybrid in own grade)	0.41 (0.40)	0.13 (0.52)	-0.01 (0.30)

Table reports coefficient on interactions of post-COVID indicator with grade and learning mode indicators; control variables include time trend and indicators for month, grade, neighborhood SES quartile, birth quarter, and any-grade and own-grade hybrid indicators. Estimation sample includes observations from September 2016 through March 2022, with sample sizes as reported in Table 3a. Standard errors clustered by student in parentheses. *** = 1%, ** = 5%, * = 10%.

Table B4i: COVID effects on physician services by learning mode, non-Indigenous English home language boys.

Description	COVID effect on monthly % receiving physician services				
	Depression/ anxiety	ADHD	Psychosis	Conduct	Substance
COVID effect w/interaction terms					
Post x (grade 8)	-0.05 (0.06)	-0.03 (0.06)	0.04 (0.03)	0.05** (0.02)	-0.03*** (0.01)
Post x (grade 9)	-0.08 (0.06)	-0.09* (0.06)	-0.01 (0.03)	0.05** (0.02)	-0.03** (0.01)
Post x (grade 10)	0.01 (0.07)	-0.07 (0.05)	-0.05 (0.03)	-0.02 (0.02)	-0.03** (0.01)
Post x (grade 11)	0.12 (0.07)	-0.03 (0.05)	-0.03 (0.03)	0.00 (0.02)	-0.03** (0.02)
Post x (hybrid in any grade)	-0.07 (0.07)	0.09 (0.07)	-0.05 (0.04)	-0.05** (0.02)	0.01 (0.01)
Post x (hybrid in own grade)	-0.09 (0.07)	-0.01 (0.07)	0.09** (0.04)	0.10*** (0.02)	0.00 (0.01)

Table reports coefficient on interactions of post-COVID indicator with grade and learning mode indicators; control variables include time trend and indicators for month, grade, neighborhood SES quartile, birth quarter, and any-grade and own-grade hybrid indicators. Estimation sample includes observations from September 2016 through March 2022, with sample sizes as reported in Table 3a. Standard errors clustered by student in parentheses. *** = 1%, ** = 5%, * = 10%.

Table B4j: COVID effects on prescriptions filled by learning mode, non-Indigenous English home language boys.

Description	COVID effect on monthly % filling prescriptions		
	Antidepressant and related	ADHD-related	Antipsychotic
COVID effect w/interaction terms:			
Post x (grade 8)	0.03 (0.10)	0.03 (0.13)	0.15** (0.06)
Post x (grade 9)	-0.04 (0.10)	-0.07 (0.11)	0.08 (0.06)
Post x (grade 10)	0.03 (0.10)	-0.17* (0.10)	0.00 (0.06)
Post x (grade 11)	0.17* (0.10)	-0.06 (0.10)	0.03 (0.06)
Post x (hybrid in any grade)	-0.21* (0.11)	-0.22 (0.14)	-0.19*** (0.07)
Post x (hybrid in own grade)	0.02 (0.12)	0.04 (0.14)	0.18** (0.07)

Table reports coefficient on interactions of post-COVID indicator with grade and learning mode indicators; control variables include time trend and indicators for month, grade, neighborhood SES quartile, birth quarter, and any-grade and own-grade hybrid indicators. Estimation sample includes observations from September 2016 through March 2022, with sample sizes as reported in Table 3a. Standard errors clustered by student in parentheses. *** = 1%, ** = 5%, * = 10%.

Table B4k: COVID effects on physician services by learning mode, non-Indigenous non-English home language boys.

Description	COVID effect on monthly % receiving physician services				
	Depression/ anxiety	ADHD	Psychosis	Conduct	Substance
COVID effect w/interaction terms					
Post x (grade 8)	0.19* (0.11)	0.12 (0.10)	-0.10 (0.08)	0.00 (0.04)	-0.01 (0.02)
Post x (grade 9)	0.20* (0.11)	0.05 (0.08)	-0.10 (0.08)	-0.01 (0.03)	0.00 (0.02)
Post x (grade 10)	0.25** (0.11)	0.03 (0.09)	-0.10 (0.08)	0.00 (0.03)	-0.02 (0.02)
Post x (grade 11)	0.30** (0.12)	0.02 (0.09)	-0.01 (0.08)	-0.01 (0.03)	-0.02 (0.02)
Post x (hybrid in any grade)	-0.18* (0.11)	-0.08 (0.09)	0.11 (0.09)	0.01 (0.04)	0.01 (0.02)
Post x (hybrid in own grade)	-0.04 (0.07)	0.04 (0.07)	-0.03 (0.04)	0.01 (0.02)	0.01 (0.01)

Table reports coefficient on interactions of post-COVID indicator with grade and learning mode indicators; control variables include time trend and indicators for month, grade, neighborhood SES quartile, birth quarter, and any-grade and own-grade hybrid indicators. Estimation sample includes observations from September 2016 through March 2022, with sample sizes as reported in Table 3a. Standard errors clustered by student in parentheses. *** = 1%, ** = 5%, * = 10%.

Table B4l: COVID effects on prescriptions filled by learning mode, non-Indigenous non-English home language boys.

Description	COVID effect on monthly % filling prescriptions		
	Antidepressant and related	ADHD-related	Antipsychotic
COVID effect w/interaction terms			
Post x (grade 8)	0.17 (0.15)	-0.08 (0.19)	-0.07 (0.11)
Post x (grade 9)	0.13 (0.15)	-0.08 (0.18)	-0.07 (0.11)
Post x (grade 10)	0.15 (0.16)	-0.02 (0.17)	-0.07 (0.11)
Post x (grade 11)	0.16 (0.16)	-0.03 (0.17)	0.03 (0.11)
Post x (hybrid in any grade)	-0.18 (0.15)	0.01 (0.19)	0.02 (0.11)
Post x (hybrid in own grade)	0.09 (0.08)	-0.05 (0.09)	-0.01 (0.05)

Table reports coefficient on interactions of post-COVID indicator with grade and learning mode indicators; control variables include time trend and indicators for month, grade, neighborhood SES quartile, birth quarter, and any-grade and own-grade hybrid indicators. Estimation sample includes observations from September 2016 through March 2022, with sample sizes as reported in Table 3a. Standard errors clustered by student in parentheses. *** = 1%, ** = 5%, * = 10%.